

**BERESKIN & PARR**

**United States**

**Title:**           **Methods and Compositions for Modulating Fertility**

**Inventors:**     **Reginald M. Gorczynski and David A. Clark**

B&P File No. 9579-034/MG

**Title: Methods and Compositions for Modulating Fertility**

5 This application is a continuation-in-part of U.S. application serial no. 09/570,367 filed May 5, 1998 (now allowed) which is a continuation of PCT/CA98/01038 filed November 6, 1998 (which designated the U.S.) which claims the benefit of U.S. serial no. 60/064,764 filed November 7, 1997 (now abandoned).

**FIELD OF THE INVENTION**

10 The present invention relates to methods and compositions for modulating an immune response. The invention includes the use of the protein OX-2 to suppress an immune response and to prevent fetal loss.

**BACKGROUND OF THE INVENTION**

15 The immune system protects the body from infectious agents and disease and is critical to our survival. However, in certain instances, the immune system can be the cause of illness. One example is in autoimmune disease wherein the immune system attacks its own host tissues, in many instances causing debilitating illness and sometimes resulting in death. Examples of autoimmune diseases include multiple sclerosis, type 1 insulin-dependent diabetes mellitus, lupus erythematosus and arthritis. A second  
20 example where the immune system can cause illness is during tissue or organ transplantation. Except in the cases of genetically identical animals, such as monozygotic twins, tissue and organ transplants are rejected by the recipient's immune system as foreign. The immune reaction against transplants is even more pronounced in transplantation across species or xenotransplantation. A third example where the immune system harms the  
25 host is during an allergic reaction where the immune system is activated by a generally innocuous antigen causing inflammation and in some cases tissue damage.

30 In order to inhibit the detrimental immune reactions during transplantation, autoimmune disease and allergic reactions, immunosuppressive drugs (such as cyclosporin A, tacrolimas, and corticosteroids) or antibody therapies (such as anti-T cell antibodies) are generally administered. Unfortunately, these non-specific modes of

immunosuppression generally have undesirable side effects. For example, cyclosporin may cause decreased renal function, hypertension, toxicity and it must be administered for the life of the patient. Corticosteroids may cause decreased resistance to infection, painful arthritis, osteoporosis and cataracts.

5 The anti-T cell antibodies may cause fever, hypertension, diarrhea or sterile meningitis and are quite expensive.

In view of the problems associated with immunosuppression, there has been an interest in developing methods or therapies that induce unresponsiveness or tolerance in the host to a transplant, to "self" tissues in  
10 autoimmune disease and to harmless antigens associated with allergies. The inventor has been studying the mechanisms involved in transplant rejection and has developed methods for inducing a state of antigen-specific immunological tolerance in transplantation. In particular, in animal allograft models, the inventor has demonstrated that graft survival can be increased  
15 if the recipient animal is given a pre-transplant infusion via the portal vein of irradiated spleen cells from the donor animal. In contrast, a pre-transplant infusion via the tail vein does not prolong graft survival.

Understanding the molecular mechanisms involved in the induction of tolerance following portal-venous (pv) immunization may lead to  
20 the development of methods of inducing immune tolerance that may be useful in transplant, autoimmune disease and allergies.

#### **SUMMARY OF THE INVENTION**

Genes that show an increase in expression following portal venous immunization have been identified. One of the genes isolated  
25 encodes OX-2, a molecule with previously unknown function belonging to the Ig superfamily. The OX-2 molecule is also generally referred to as CD200 in the current literature. The inventors have shown that administering antibodies to OX-2 inhibited the graft survival generally seen following pre-transplant pv immunization. The inventors have also shown that there is a negative  
30 association between levels of OX-2 and risk of fetal loss. In particular, the inventors have shown administering OX-2 reduced fetal loss rates while inhibiting OX-2 reversed the effect. The inventors have further shown that

OX-2 inhibits cytotoxic cells and IL-2 production and induces IL-4 production. All of these results demonstrate that OX-2 is involved in immune suppression.

Consequently, broadly stated, the present invention provides a method of suppressing an immune system comprising administering an effective amount of an OX-2 protein or a nucleic acid sequence encoding an OX-2 protein to an animal in need of such treatment.

In one embodiment, the present invention provides a method of preventing or inhibiting fetal loss comprising administering an effective amount of an OX-2 protein or a nucleic acid sequence encoding an OX-2 protein to an animal in need thereof.

The invention also includes pharmaceutical compositions containing OX-2 proteins or nucleic acids encoding OX-2 proteins for use in inducing tolerance in transplantation or autoimmune disease.

As stated above, OX-2 can be used to induce immune suppression. Consequently, inhibiting OX-2 may also be useful in preventing immune suppression.

Therefore, in another aspect, the present invention provides a method of preventing immune suppression comprising administering an effective amount of an agent that inhibits OX-2 to an animal in need thereof. In a preferred embodiment the OX-2 inhibitor is an antibody that binds OX-2 or an antisense oligonucleotide that inhibits the expression of OX-2.

In one embodiment, the present invention provides a method of inducing fetal loss comprising administering an effective amount of an agent that inhibits OX-2 to an animal in need thereof.

The invention also includes pharmaceutical compositions containing an OX-2 inhibitor for use in inducing or augmenting an immune response.

Other features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples while indicating preferred embodiments of the invention are given by way of illustration only, since various changes and modifications within the spirit and



scope of the invention will become apparent to those skilled in the art from this detailed description.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

The invention will now be described in relation to the drawings in which:

Figure 1 illustrates PCR validation of suppressive subtractive hybridization using  $\beta$ -actin primers.

Figure 2 illustrates PCR validation of suppressive subtractive hybridization using IL-10 primers.

Figure 3 is an autoradiograph using  $^{32}\text{P}$ -labeled probes from 4 clones obtained from the subtractive hybridization process.

Figure 4 is flow cytometry profile of spleen adherent cells.

Figure 5A and B are Western Blots illustrating the increased expression of OX-2 antigen after pv immunization. Figure 5A shows staining with a control mouse antibody, anti-mouse CD8a. Figure 5B shows staining with anti-rat MRC OX-2.

Figure 6 is a graft showing percent survival versus days post renal transplantation.

Figure 7 shows the cDNA sequence of rat (SEQ.ID.NO.:20), mouse (SEQ.ID.NO.:22) and human MRC OX-2 (SEQ.ID.NO.:18).

Figure 8 shows the deduced protein sequence of rat (SEQ.ID.NO.:21), mouse (SEQ.ID.NO.:2) and human MRC OX-2 (SEQ.ID.NO.:19) protein.

Figures 9A and 9B are bar graphs showing cytokine production and cell proliferation following stimulation by allogeneic DC using hepatic NPMC.

Figures 10A, 10B and 10C are bar graphs showing inhibition of cell proliferation and cytokine production by hepatic NPMC.

Figure 11 is a bar graph analysis of FACS data showing OX-2 expression in a subpopulation of NPC.

Figure 12 shows PCR analysis mRNA expression of B7-1, B7-2 and OX-2 in various hepatic NPMC cell fractions.

Figures 13A and 13B are bar graphs showing proliferation and cytokine production by NPMC from *Flt3L* treated mice.

Figure 14 is a bar graph showing cytokines produced from C3H mice with C57BL16 renal allografts and NPC from *Flt3* treated C57BL16 donors.

Figure 15 is a graph showing inhibition of graft rejection with NPC from *Flt3* treated mice.

Figure 16 is a graph showing that anti-OX-2 reverses inhibition by NPC. The effect of anti-B7-1, anti-B7-2 and anti-OX-2 on primary allostimulation is shown.

Figure 17 is a graph showing that anti-OX-2 mAb reverses inhibition by NPC and inhibits the development of immunoregulatory cells.

Figure 18A is a photograph showing *in situ* hybridization with antisense OX-2 in a day 8.5 implantation site from a mouse CBA/JxDBA/2 that is susceptible to fetal loss.

Figure 18B is a photograph showing *in situ* hybridization with antisense OX-2 in a day 8.5 implantation site from a mouse DBA/2XCBA/J that is not susceptible to spontaneous fetal loss.

Figure 19 is a graph showing the effect of anti-OX-2 monoclonal antibody 3B6 on spontaneous resorption (abortion) rates. Error bars show 1 standard deviation. A minimum of 20 implants was scored for each group. \* significant increase in abortion rate,  $P < 0.02$ .

Figure 20 is a graph showing effect of OX2 immunoadhesion (OX2:Fc) on spontaneous resorption (abortion) rates or renal allograft rejection. Error bars show 1 standard deviation. A minimum of 45 implants was scored for each group of pregnant mice (mice received a single dose of OX2:Fc at day 6.5; 10 mice/group were used for renal allografts, and mice received multiple iv injections of OX2:Fc, beginning on the day of transplant, and at 2 day intervals thereafter. \* significant decrease in abortion rate, or increased graft survival, relative to controls.

Figure 21 shows the importance of CD200 (OX-2) in rescuing potentially doomed embryos which are low in OX-2 mRNA expression by the *in situ* hybridisation result.

Figure 22 shows the molecule size ladder (lane 1), trophoblast with full length primers (lane 2) and exon 3 primers (lane 3). The full length and shorter mRNA OX-2 transcripts are seen. Lane 4 (corresponding to lane 2) and lane 5 (corresponding to lane 3) represent negative result obtained with CD9<sup>+</sup> stromal cells.

Figure 23 shows result using antibody detecting CK5,6,8,17; similar results have been obtained using FITC-anti-CK18. The upper left panel shows the forward and side scatter pattern. FITC and PE isotype controls used to set gates is shown below on left. Upper right panel shows cytokeratin-positive cells, the majority of which were also OX-2<sup>+</sup>. A significant population of CK- OX-2<sup>++</sup> cells was also noted. The identity of this population has not been established. On the right, the middle panel shows staining profile for OX-2 for the CK- and CK<sup>+</sup> populations, and the lower panel shows the scatter. Clearly these populations have different properties. Velocity sedimentation separation indicates both populations may have OX-2-dependent immunoregulatory activity.

#### **DETAILED DESCRIPTION OF THE INVENTION**

The present inventors have identified genes that show an increase in expression following portal venous immunization. These genes play a role in the development of immune suppression or tolerance and may be useful in developing therapies for the prevention and treatment of transplant rejection, fetal loss, autoimmune disease or allergies.

Using suppression subtractive hybridization (SSH), the inventor has isolated a clone that is preferentially expressed in mice receiving allogenic renal grafts along with pre-transplant donor-specific immunization and that encodes the protein OX-2. The OX-2 protein (also known as MRC OX-2) in rat was described as a 41Kd-47Kd glycoprotein which is expressed on the cell surface of thymocytes, follicular dendritic cells and endothelium, B cells and neuronal cells. Differences in apparent size of the molecule in different tissues is probably a function of differential glycosylation. The function of the molecule was previously unknown, but DNA and amino acid sequence analysis shows it has a high degree of homology to molecules of the immunoglobulin gene family, which includes molecules important in

lymphocyte antigen recognition and cell-cell interaction (e.g. CD4, CD8, ICAMs, VCAMs), as well as adhesion receptor molecules (NCAMs) in the nervous system. Members of the immunoglobulin superfamily are distinct from other molecules of the integrin and selectin families, which, at least within the immune system, also seem to play critical role in cell recognition, migration and even development of the lymphocyte recognition repertoire (by regulating intra-thymic selection events). It has become increasingly evident that molecules of these different families play an important role in human disease.

The inventor has shown that administering antibodies to OX-2 inhibited the graft survival generally seen following pre-transplant pv immunization. The inventor has also shown that there is negative association between levels of OX-2 and risk of fetal loss and that administering OX-2 prevents fetal loss and inhibiting OX-2 causes fetal loss. The inventor has further shown that OX-2 inhibits cytotoxic cells and IL-2 production and induces IL-4 production. The data supports the role of OX-2 in immune suppression.

## **THERAPEUTIC METHODS**

### ***Inducing Immune Suppression***

In one aspect, the present invention provides a method of suppressing an immune response comprising administering an effective amount of an OX-2 protein or a nucleic acid sequence encoding an OX-2 protein to an animal in need of such treatment. The invention includes a use of an effective amount of an OX-2 protein or a nucleic acid sequence encoding an OX-2 protein to suppress an immune response.

The term "OX-2 protein" includes OX-2 or CD200 from any species or source and includes a full length OX-2 protein as well as fragments or portions of the protein. The term "OX-2" is also generally referred to as "CD200" due to a change in nomenclature. Both "OX-2" and "CD200" may be used interchangeably in the application. Preferred fragments or portions of the OX-2 or CD200 protein are those that are sufficient to suppress an immune response. Determining whether a particular OX-2 or CD200 protein can suppress an immune response can be assessing using known *in vitro*

immune assays including, but not limited to, inhibiting a mixed leucocyte reaction; inhibiting a cytotoxic T cell response; inhibiting interleukin-2 production; inhibiting IFN $\gamma$  production; inhibiting a Th1 cytokine profile; inducing IL-4 production; inducing TGF $\beta$  production; inducing IL-10 production; inducing a Th2 cytokine profile; and any other assay that would be known to one of skill in the art to be useful in detecting immune suppression.

One of skill in the art can also determine whether or not a particular OX-2 or fragment thereof is useful in preventing fetal loss. As mentioned above, one of skill in the art can readily test an OX-2 or OX-2 fragment for its ability to suppress an immune response using known in vitro assays. In addition the OX-2 or OX-2 fragment can also be tested for its ability to prevent fetal loss in an animal model. For example, one could use the model described in Example 8 wherein the ability of OX-2 to prevent cytokine induced abortion in abortion-prone CBA x DBA/2 mice is assessed. Further, mice pre-immunized with anti-phospholipid may also be used.

The term "administering an OX-2 protein" includes both the administration of the OX-2 protein as well as the administration of a nucleic acid sequence encoding an OX-2 protein. In the latter case, the OX-2 protein is produced in vivo in the animal.

In a preferred embodiment, the OX-2 protein is prepared and administered as a soluble fusion protein. The fusion protein may contain the extracellular domain of OX-2 linked to an immunoglobulin (Ig) Fc Region. The OX-2 fusion may be prepared using techniques known in the art. Generally, a DNA sequence encoding the extracellular domain of OX-2 is linked to a DNA sequence encoding the Fc of the Ig and expressed in an appropriate expression system where the OX-2 - FcIg fusion protein is produced.

The OX-2 or protein may be obtained from known sources or prepared using recombinant DNA techniques. The protein may have any of the known published sequences for OX-2 or CD200. The sequences can be obtained from GenBank. The human sequence has accession no. M17226 X0523; the rat sequence has accession no. X01785; and the mouse sequence has accession no. AF029214. The nucleic acid and protein sequences of OX-2 (CD200) from human, mouse and rat are also shown in

Figures 7 and 8 and in SEQ.ID.Nos.: 18, 22 and 20 (nucleic acid) and SEQ.ID.Nos.:19, 21 and 2 (protein).

5 The OX-2 protein may also be modified to contain amino acid substitutions, insertions and/or deletions that do not alter the immunosuppressive properties of the protein. Conserved amino acid substitutions involve replacing one or more amino acids of the OX-2 amino acid sequence with amino acids of similar charge, size, and/or hydrophobicity characteristics. When only conserved substitutions are made the resulting analog should be functionally equivalent to the OX-2 protein. Non-conserved  
10 substitutions involve replacing one or more amino acids of the OX-2 amino acid sequence with one or more amino acids which possess dissimilar charge, size, and/or hydrophobicity characteristics.

The OX-2 protein may be modified to make it more therapeutically effective or suitable. For example, the OX-2 protein may be  
15 cyclized as cyclization allows a peptide to assume a more favourable conformation. Cyclization of the OX-2 peptides may be achieved using techniques known in the art. In particular, disulphide bonds may be formed between two appropriately spaced components having free sulfhydryl groups. The bonds may be formed between side chains of amino acids, non-amino  
20 acid components or a combination of the two. In addition, the OX-2 protein or peptides of the present invention may be converted into pharmaceutical salts by reacting with inorganic acids including hydrochloric acid, sulphuric acid, hydrobromic acid, phosphoric acid, etc., or organic acids including formic acid, acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic  
25 acid, succinic acid, malic acid, tartaric acid, citric acid, benzoic acid, salicylic acid, benzenesulphonic acid, and tolunesulphonic acids.

Administration of an "effective amount" of the OX-2 protein and nucleic acid of the present invention is defined as an amount effective, at dosages and for periods of time necessary to achieve the desired result. The  
30 effective amount of the OX-2 protein or nucleic acid of the invention may vary according to factors such as the disease state, age, sex, and weight of the animal. Dosage regima may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily or

the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation.

The term "animal" as used herein includes all members of the animal kingdom including humans. When treating fetal loss, the animal is a female who is desirous of becoming pregnant or maintaining a pregnancy.

The present inventors have shown that there is an association between levels of OX-2 expression and fertility. In particular the inventor has shown that low levels (or no levels) of OX-2 is related to fetal loss. Further, administering a OX-2:Fc fusion protein prevented fetal loss.

Accordingly, the present invention provides a method of preventing, reducing or inhibiting fetal loss comprising administering an effective amount of an OX-2 protein or a nucleic acid sequence encoding an OX-2 protein to an animal in need thereof. The invention includes a use of an effective amount of an OX-2 protein on a nucleic acid molecules encoding an OX-2 protein to prevent or inhibit fetal loss. The OX-2 protein may be from any species and may be the full length sequence or a fragment thereof that is capable of preventing or inhibiting fetal loss.

In another embodiment, the present invention provides a method of inducing immune tolerance to a transplanted organ or tissue in a recipient animal comprising administering an effective amount of a OX-2 protein or a nucleic acid sequence encoding an OX-2 protein to the recipient animal prior to the transplantation of the organ or tissue. The invention includes a use of an effective amount of a OX-2 protein or a nucleic acid sequence encoding an OX-2 protein to induce immune tolerance to a transplanted organ or tissue.

The term "inducing immune tolerance" means rendering the immune system unresponsive to a particular antigen without inducing a prolonged generalized immune deficiency. The term "antigen" means a substance that is capable of inducing an immune response. In the case of autoimmune disease, immune tolerance means rendering the immune system unresponsive to an auto-antigen that the host is recognizing as foreign, thus causing an autoimmune response. In the case of allergy, immune tolerance means rendering the immune system unresponsive to an allergen that

generally causes an immune response in the host. In the case of transplantation, immune tolerance means rendering the immune system unresponsive to the antigens on the transplant. An alloantigen refers to an antigen found only in some members of a species, such as blood group antigens. A xenoantigen refers to an antigen that is present in members of one species but not members of another. Correspondingly, an allograft is a graft between members of the same species and a xenograft is a graft between members of a different species.

The recipient can be any member of the animal kingdom including rodents, pigs, cats, dogs, ruminants, non-human primates and preferably humans. The organ or tissue to be transplanted can be from the same species as the recipient (allograft) or can be from another species (xenograft). The tissues or organs can be any tissue or organ including heart, liver, kidney, lung, pancreas, pancreatic islets, brain tissue, cornea, bone, intestine, skin and hematopoietic cells.

The method of the invention may be used to prevent graft versus host disease wherein the immune cells in the transplant mount an immune attack on the recipient's immune system. This can occur when the tissue to be transplanted contains immune cells such as when bone marrow or lymphoid tissue is transplanted when treating leukemias, aplastic anemias and enzyme or immune deficiencies, for example.

Accordingly, in another embodiment, the present invention provides a method of preventing or inhibiting graft versus host disease in a recipient animal receiving an organ or tissue transplant comprising administering an effective amount of a OX-2 protein or a nucleic acid sequence encoding an OX-2 protein to the organ or tissue prior to the transplantation in the recipient animal. The invention includes a use of an effective amount of an OX-2 protein or a nucleic acid molecule encoding an OX-2 protein to prevent or inhibit graft versus host disease.

As stated previously, the method of the present invention may also be used to treat or prevent autoimmune disease. In an autoimmune disease, the immune system of the host fails to recognize a particular antigen as "self" and an immune reaction is mounted against the host's tissues



expressing the antigen. Normally, the immune system is tolerant to its own host's tissues and autoimmunity can be thought of as a breakdown in the immune tolerance system.

Accordingly, in a further embodiment, the present invention provides a method of preventing or treating an autoimmune disease comprising administering an effective amount of an OX-2 protein or a nucleic acid sequence encoding an OX-2 protein to an animal having, suspected of having, or susceptible to having an autoimmune disease. The invention includes a use of an effective amount of an OX-2 protein on a nucleic acid molecule encoding an OX-2 protein to prevent or inhibit an autoimmune disease.

Autoimmune diseases that may be treated or prevented according to the present invention include, but are not limited to, type 1 insulin-dependent diabetes mellitus, adult respiratory distress syndrome, inflammatory bowel disease, dermatitis, meningitis, thrombotic thrombocytopenic purpura, Sjögren's syndrome, encephalitis, uveitis, leukocyte adhesion deficiency, rheumatoid arthritis, rheumatic fever, Reiter's syndrome, psoriatic arthritis, progressive systemic sclerosis, primary biliary cirrhosis, pemphigus, pemphigoid, necrotizing vasculitis, myasthenia gravis, multiple sclerosis, lupus erythematosus, polymyositis, sarcoidosis, granulomatosis, vasculitis, pernicious anemia, CNS inflammatory disorder, antigen-antibody complex mediated diseases, autoimmune haemolytic anemia, Hashimoto's thyroiditis, Graves disease, habitual spontaneous abortions, Reynard's syndrome, glomerulonephritis, dermatomyositis, chronic active hepatitis, celiac disease, autoimmune complications of AIDS, atrophic gastritis, ankylosing spondylitis and Addison's disease.

As stated previously, the method of the present invention may also be used to treat or prevent an allergic reaction. In an allergic reaction, the immune system mounts an attack against a generally harmless, innocuous antigen or allergen. Allergies that may be prevented or treated using the methods of the invention include, but are not limited to, hay fever, asthma, atopic eczema as well as allergies to poison oak and ivy, house dust mites, bee pollen, nuts, shellfish, penicillin and numerous others.

Accordingly, in a further embodiment, the present invention provides a method of preventing or treating an allergy comprising administering an effective amount of an OX-2 protein or a nucleic acid sequence encoding an OX-2 protein to an animal having or suspected of having an allergy. The invention includes a use of an effective amount of an OX-2 protein or a nucleic acid molecule encoding an OX-2 protein to prevent or treat an allergy.

### ***Preventing Immune Suppression***

In another aspect, the present invention provides a method of preventing immune suppression comprising administering an effective amount of an agent that inhibits OX-2 to an animal in need thereof.

There are a large number of situations whereby it is desirable to prevent immune suppression including, but not limited to, the treatment of infections, cancer and Acquired Immune Deficiency Syndrome and the induction of fetal loss.

Accordingly, the present invention provides a method of preventing immune suppression comprising administering an effective amount of an agent that inhibits OX-2 to an animal in need thereof. In one embodiment the present invention provides a method of inducing fetal loss comprising administering an effective amount of an agent that inhibits OX-2 to an animal in need thereof. The animal is preferably a pregnant female.

The agent that inhibits OX-2 can be any agent that decreases the expression or activity of an OX-2 protein such that the immune suppression caused by OX-2 is reduced, inhibited and/or prevented. One of skill in the art can readily determine whether or not a particular agent is effective in inhibiting OX-2. For example, the agent can be tested in *in vitro* assays to determine if the function or activity of OX-2 is inhibited. The agent can also be tested for its ability to induce an immune response using *in vitro* immune assays including, but not limited to, enhancing a cytotoxic T cell response; inducing interleukin-2 (IL-2) production; inducing IFN $\gamma$  production; inducing a Th1 cytokine profile; inhibiting IL-4 production; inhibiting TGF $\beta$  production; inhibiting IL-10 production; inhibiting a Th2 cytokine profile and

any other assay that would be known to one of skill in the art to be useful in detecting immune activation.

One of skill in the art can also determine whether a particular agent is useful in inducing fetal loss. As mentioned above, one can test the agent for its ability to induce an immune response using known in vitro assays. In addition, the agent can be tested in an animal model, for example as described in Example 11, wherein the agent is administered to a pregnant rodent.

In a preferred embodiment, the agent that inhibits OX-2 is an OX-2 specific antibody. The present inventor has prepared antibodies to OX-2 which are described in Examples 4 and 5. Antibodies to OX-2 may also be prepared using techniques known in the art such as those described by Kohler and Milstein, Nature 256, 495 (1975) and in U.S. Patent Nos. RE 32,011; 4,902,614; 4,543,439; and 4,411,993, which are incorporated herein by reference. (See also Monoclonal Antibodies, Hybridomas: A New Dimension in Biological Analyses, Plenum Press, Kennett, McKearn, and Bechtol (eds.), 1980, and Antibodies: A Laboratory Manual, Harlow and Lane (eds.), Cold Spring Harbor Laboratory Press, 1988, which are also incorporated herein by reference). Within the context of the present invention, antibodies are understood to include monoclonal antibodies, polyclonal antibodies, antibody fragments (e.g., Fab, and F(ab')<sub>2</sub>) and recombinantly produced binding partners.

Accordingly, the present invention provides a method of preventing immune suppression or inducing fetal loss comprising administering an effective amount of an antibody that inhibits OX-2 to an animal in need thereof.

In another embodiment, the OX-2 inhibitor is an antisense oligonucleotide that inhibits the expression of OX-2. Antisense oligonucleotides that are complimentary to a nucleic acid sequence from an OX-2 gene can be used in the methods of the present invention to inhibit OX-2. The present inventor has prepared antisense oligonucleotides to OX-2 which are described in Example 3.

Accordingly, the present invention provides a method of preventing immune suppression comprising administering an effective amount of an antisense oligonucleotide that is complimentary to a nucleic acid sequence from a OX-2 gene to an animal in need thereof.

5           The term antisense oligonucleotide as used herein means a nucleotide sequence that is complimentary to its target.

          In one embodiment of the invention, the present invention provides an antisense oligonucleotide that is complimentary to a nucleic acid molecule having a sequence as shown in Figure 7, wherein T can also be U,  
10           or a fragment thereof.

          The term "oligonucleotide" refers to an oligomer or polymer of nucleotide or nucleoside monomers consisting of naturally occurring bases, sugars, and intersugar (backbone) linkages. The term also includes modified or substituted oligomers comprising non-naturally occurring monomers or portions thereof, which function similarly. Such modified or substituted  
15           oligonucleotides may be preferred over naturally occurring forms because of properties such as enhanced cellular uptake, or increased stability in the presence of nucleases. The term also includes chimeric oligonucleotides which contain two or more chemically distinct regions. For example, chimeric  
20           oligonucleotides may contain at least one region of modified nucleotides that confer beneficial properties (e.g. increased nuclease resistance, increased uptake into cells), or two or more oligonucleotides of the invention may be joined to form a chimeric oligonucleotide.

          The antisense oligonucleotides of the present invention may be  
25           ribonucleic or deoxyribonucleic acids and may contain naturally occurring bases including adenine, guanine, cytosine, thymidine and uracil. The oligonucleotides may also contain modified bases such as xanthine, hypoxanthine, 2-aminoadenine, 6-methyl, 2-propyl and other alkyl adenines, 5-halo uracil, 5-halo cytosine, 6-aza uracil, 6-aza cytosine and 6-aza thymine,  
30           pseudo uracil, 4-thiouracil, 8-halo adenine, 8-aminoadenine, 8-thiol adenine, 8-thiolalkyl adenines, 8-hydroxyl adenine and other 8-substituted adenines, 8-halo guanines, 8-amino guanine, 8-thiol guanine, 8-thiolalkyl guanines, 8-hydroxyl guanine and other 8-substituted guanines, other aza and deaza

uracils, thymidines, cytosines, adenines, or guanines, 5-trifluoromethyl uracil and 5-trifluoro cytosine.

Other antisense oligonucleotides of the invention may contain modified phosphorous, oxygen heteroatoms in the phosphate backbone, short chain alkyl or cycloalkyl intersugar linkages or short chain heteroatomic or heterocyclic intersugar linkages. For example, the antisense oligonucleotides may contain phosphorothioates, phosphotriesters, methyl phosphonates, and phosphorodithioates. In an embodiment of the invention there are phosphorothioate bonds links between the four to six 3'-terminus bases. In another embodiment phosphorothioate bonds link all the nucleotides.

The antisense oligonucleotides of the invention may also comprise nucleotide analogs that may be better suited as therapeutic or experimental reagents. An example of an oligonucleotide analogue is a peptide nucleic acid (PNA) wherein the deoxyribose (or ribose) phosphate backbone in the DNA (or RNA), is replaced with a polyamide backbone which is similar to that found in peptides (P.E. Nielsen, et al Science 1991, 254, 1497). PNA analogues have been shown to be resistant to degradation by enzymes and to have extended lives *in vivo* and *in vitro*. PNAs also bind stronger to a complimentary DNA sequence due to the lack of charge repulsion between the PNA strand and the DNA strand. Other oligonucleotides may contain nucleotides containing polymer backbones, cyclic backbones, or acyclic backbones. For example, the nucleotides may have morpholino backbone structures (U.S. Pat. No. 5,034,506). Oligonucleotides may also contain groups such as reporter groups, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an antisense oligonucleotide. Antisense oligonucleotides may also have sugar mimetics.

The antisense nucleic acid molecules may be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. The antisense nucleic acid molecules of the invention or a fragment thereof, may be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the

biological stability of the molecules or to increase the physical stability of the duplex formed with mRNA or the native gene e.g. phosphorothioate derivatives and acridine substituted nucleotides. The antisense sequences may be produced biologically using an expression vector introduced into cells in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense sequences are produced under the control of a high efficiency regulatory region, the activity of which may be determined by the cell type into which the vector is introduced.

### **COMPOSITIONS**

The invention also includes pharmaceutical compositions containing OX-2 proteins or nucleic acids for use in immune suppression as well as pharmaceutical compositions containing an OX-2 inhibitor for use in preventing immune suppression.

Such pharmaceutical compositions can be for intralesional, intravenous, topical, rectal, parenteral, local, inhalant or subcutaneous, intradermal, intramuscular, intrathecal, transperitoneal, oral, and intracerebral use. The composition can be in liquid, solid or semisolid form, for example pills, tablets, creams, gelatin capsules, capsules, suppositories, soft gelatin capsules, gels, membranes, tubelets, solutions or suspensions.

The pharmaceutical compositions of the invention can be intended for administration to humans or animals. Dosages to be administered depend on individual needs, on the desired effect and on the chosen route of administration.

The pharmaceutical compositions can be prepared by per se known methods for the preparation of pharmaceutically acceptable compositions which can be administered to patients, and such that an effective quantity of the active substance is combined in a mixture with a pharmaceutically acceptable vehicle. Suitable vehicles are described, for example, in Remington's Pharmaceutical Sciences (Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., USA 1985).

On this basis, the pharmaceutical compositions include, albeit not exclusively, the active compound or substance in association with one or

more pharmaceutically acceptable vehicles or diluents, and contained in buffered solutions with a suitable pH and iso-osmotic with the physiological fluids. The pharmaceutical compositions may additionally contain other agents such as immunosuppressive drugs or antibodies to enhance immune tolerance or immunostimulatory agents to enhance the immune response.

In one embodiment, the pharmaceutical composition for use in preventing fetal loss comprises an effective amount of a OX-2 protein in admixture with a pharmaceutically acceptable diluent or carrier. The OX-2 protein is preferably prepared as an immunoadhesion molecule in soluble form which can be administered to the patient.

In another embodiment, the pharmaceutical composition for use in preventing fetal loss comprises an effective amount of a nucleic acid molecule encoding a OX-2 protein in admixture with a pharmaceutically acceptable diluent or carrier.

The nucleic acid molecules of the invention encoding a OX-2 protein may be used in gene therapy to induce immune tolerance. Recombinant molecules comprising a nucleic acid sequence encoding a OX-2 protein, or fragment thereof, may be directly introduced into cells or tissues *in vivo* using delivery vehicles such as retroviral vectors, adenoviral vectors and DNA virus vectors. They may also be introduced into cells *in vivo* using physical techniques such as microinjection and electroporation or chemical methods such as coprecipitation and incorporation of DNA into liposomes. Recombinant molecules may also be delivered in the form of an aerosol or by lavage. The nucleic acid molecules of the invention may also be applied extracellularly such as by direct injection into cells. The nucleic acid molecules encoding OX-2 are preferably prepared as a fusion with a nucleic acid molecule encoding an immunoglobulin (Ig) Fc region. As such, the OX-2 protein will be expressed *in vivo* as a soluble fusion protein.

In another aspect, the pharmaceutical composition for use in inducing fetal loss comprises an effective amount of a OX-2 inhibitor in admixture with a pharmaceutically acceptable diluent or carrier. Such compositions may be administered as a vaccine either alone or in combination with other active agents.

In one embodiment, the pharmaceutical composition for use in inducing fetal loss comprises an effective amount of an antibody to OX-2 in admixture with a pharmaceutically acceptable diluent or carrier. The antibodies may be delivered intravenously.

5 In another embodiment, the pharmaceutical composition for use in inducing fetal loss comprises an effective amount of an antisense oligonucleotide nucleic acid complimentary to a nucleic acid sequence from a OX-2 gene in admixture with a pharmaceutically acceptable diluent or carrier. The oligonucleotide molecules may be administered as described above for  
10 the compositions containing OX-2 nucleic acid sequences.

#### **MURINE OX-2**

The inventor has cloned and sequenced the murine OX-2 gene. Accordingly, the invention also includes an isolated nucleic acid sequence encoding a murine OX-2 gene and having the sequence shown in Figure 7  
15 and SEQ.ID.NO.:1.

The term "isolated" refers to a nucleic acid substantially free of cellular material or culture medium when produced by recombinant DNA techniques, or chemical precursors, or other chemicals when chemically synthesized. The term "nucleic acid" is intended to include DNA and RNA  
20 and can be either double stranded or single stranded.

Preferably, the purified and isolated nucleic acid molecule of the invention comprises (a) a nucleic acid sequence as shown in SEQ.ID.NO.:1, wherein T can also be U; (b) nucleic acid sequences complementary to (a); (c) a fragment of (a) or (b) that is at least 15 bases, preferably 20 to 30 bases,  
25 and which will hybridize to (a) or (b) under stringent hybridization conditions; or (a) a nucleic acid molecule differing from any of the nucleic acids of (a) or (b) in codon sequences due to the degeneracy of the genetic code.

It will be appreciated that the invention includes nucleic acid molecules encoding truncations of the murine OX-2 proteins of the invention, and analogs and homologs of the proteins of the invention and truncations  
30 thereof, as described below. It will further be appreciated that variant forms of the nucleic acid molecules of the invention which arise by alternative splicing



of an mRNA corresponding to a cDNA of the invention are encompassed by the invention.

An isolated nucleic acid molecule of the invention which is DNA can also be isolated by selectively amplifying a nucleic acid encoding a novel protein of the invention using the polymerase chain reaction (PCR) methods and cDNA or genomic DNA. It is possible to design synthetic oligonucleotide primers from the nucleic acid molecules as shown in Figure 7 for use in PCR. A nucleic acid can be amplified from cDNA or genomic DNA using these oligonucleotide primers and standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. It will be appreciated that cDNA may be prepared from mRNA, by isolating total cellular mRNA by a variety of techniques, for example, by using the guanidinium-thiocyanate extraction procedure of Chirgwin et al., Biochemistry, 18, 5294-5299 (1979). cDNA is then synthesized from the mRNA using reverse transcriptase (for example, Moloney MLV reverse transcriptase available from Gibco/BRL, Bethesda, MD, or AMV reverse transcriptase available from Seikagaku America, Inc., St. Petersburg, FL).

An isolated nucleic acid molecule of the invention which is RNA can be isolated by cloning a cDNA encoding a novel protein of the invention into an appropriate vector which allows for transcription of the cDNA to produce an RNA molecule which encodes a OX-2 protein of the invention. For example, a cDNA can be cloned downstream of a bacteriophage promoter, (e.g. a T7 promoter) in a vector, cDNA can be transcribed *in vitro* with T7 polymerase, and the resultant RNA can be isolated by standard techniques.

A nucleic acid molecule of the invention may also be chemically synthesized using standard techniques. Various methods of chemically synthesizing polydeoxynucleotides are known, including solid-phase synthesis which, like peptide synthesis, has been fully automated in commercially available DNA synthesizers (See e.g., Itakura et al. U.S. Patent No. 4,598,049; Caruthers et al. U.S. Patent No. 4,458,066; and Itakura U.S. Patent Nos. 4,401,796 and 4,373,071).

The sequence of a nucleic acid molecule of the invention may be inverted relative to its normal presentation for transcription to produce an antisense nucleic acid molecule. Preferably, an antisense sequence is constructed by inverting a region preceding the initiation codon or an  
5 unconserved region. In particular, the nucleic acid sequences contained in the nucleic acid molecules of the invention or a fragment thereof, preferably a nucleic acid sequence shown in Figure 7 may be inverted relative to its normal presentation for transcription to produce antisense nucleic acid molecules.

10 The antisense nucleic acid molecules of the invention or a fragment thereof, may be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed with mRNA or the native gene e.g. phosphorothioate  
15 derivatives and acridine substituted nucleotides. The antisense sequences may be produced biologically using an expression vector introduced into cells in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense sequences are produced under the control of a high efficiency regulatory region, the activity of which may be determined by the cell type into  
20 which the vector is introduced.

The invention also provides nucleic acids encoding fusion proteins comprising a OX-2 protein of the invention and a selected protein, or a selectable marker protein.

25 The invention further includes an isolated protein which has the amino acid sequence as shown in Figure 8 and SEQ.ID.NO.:2.

30 Within the context of the present invention, a protein of the invention may include various structural forms of the primary protein which retain biological activity. For example, a protein of the invention may be in the form of acidic or basic salts or in neutral form. In addition, individual amino acid residues may be modified by oxidation or reduction.

In addition to the full length amino acid sequence (Figure 8), the protein of the present invention may also include truncations of the protein, and analogs, and homologs of the protein and truncations thereof as

described herein. Truncated proteins may comprise peptides of at least fifteen amino acid residues.

5           Analog of the protein having the amino acid sequence shown in Figure 8, and/or truncations thereof as described herein, may include, but are not limited to an amino acid sequence containing one or more amino acid substitutions, insertions, and/or deletions. Amino acid substitutions may be of a conserved or non-conserved nature. Conserved amino acid substitutions involve replacing one or more amino acids of the proteins of the invention with amino acids of similar charge, size, and/or hydrophobicity characteristics. 10       When only conserved substitutions are made the resulting analog should be functionally equivalent. Non-conserved substitutions involve replacing one or more amino acids of the amino acid sequence with one or more amino acids which possess dissimilar charge, size, and/or hydrophobicity characteristics.

15           One or more amino acid insertions may be introduced into the amino acid sequences shown in Figure 8. Amino acid insertions may consist of single amino acid residues or sequential amino acids ranging from 2 to 15 amino acids in length. For example, amino acid insertions may be used to render the protein is no longer active. This procedure may be used *in vivo* to inhibit the activity of a protein of the invention.

20           Deletions may consist of the removal of one or more amino acids, or discrete portions from the amino acid sequence shown in Figures 8. The deleted amino acids may or may not be contiguous. The lower limit length of the resulting analog with a deletion mutation is about 10 amino acids, preferably 100 amino acids.

25           Analog of a protein of the invention may be prepared by introducing mutations in the nucleotide sequence encoding the protein. Mutations in nucleotide sequences constructed for expression of analogs of a protein of the invention must preserve the reading frame of the coding sequences. Furthermore, the mutations will preferably not create 30       complementary regions that could hybridize to produce secondary mRNA structures, such as loops or hairpins, which could adversely affect translation of the receptor mRNA.

Mutations may be introduced at particular loci by synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling ligation to fragments of the native sequence. Following ligation, the resulting reconstructed sequence encodes an analog having the desired amino acid insertion, substitution, or deletion.

Alternatively, oligonucleotide-directed site specific mutagenesis procedures may be employed to provide an altered gene having particular codons altered according to the substitution, deletion, or insertion required. Deletion or truncation of a protein of the invention may also be constructed by utilizing convenient restriction endonuclease sites adjacent to the desired deletion. Subsequent to restriction, overhangs may be filled in, and the DNA religated. Exemplary methods of making the alterations set forth above are disclosed by Sambrook et al (Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, 1989).

The invention also contemplates isoforms of the proteins of the invention. An isoform contains the same number and kinds of amino acids as a protein of the invention, but the isoform has a different molecular structure. The isoforms contemplated by the present invention are those having the same properties as a protein of the invention as described herein.

The present invention also includes a protein of the invention conjugated with a selected protein, or a selectable marker protein to produce fusion proteins. Additionally, immunogenic portions of a protein of the invention are within the scope of the invention.

The proteins of the invention (including truncations, analogs, etc.) may be prepared using recombinant DNA methods. Accordingly, the nucleic acid molecules of the present invention having a sequence which encodes a protein of the invention may be incorporated in a known manner into an appropriate expression vector which ensures good expression of the protein. Possible expression vectors include but are not limited to cosmids, plasmids, or modified viruses (e.g. replication defective retroviruses, adenoviruses and adeno-associated viruses), so long as the vector is compatible with the host cell used. The expression vectors are "suitable for transformation of a host cell", means that the expression vectors contain a

nucleic acid molecule of the invention and regulatory sequences selected on the basis of the host cells to be used for expression, which is operatively linked to the nucleic acid molecule. Operatively linked is intended to mean that the nucleic acid is linked to regulatory sequences in a manner which allows expression of the nucleic acid.

The invention therefore contemplates a recombinant expression vector of the invention containing a nucleic acid molecule of the invention, or a fragment thereof, and the necessary regulatory sequences for the transcription and translation of the inserted protein-sequence. Such expression vectors may be useful in the above-described therapies using a nucleic acid sequence encoding a OX-2 protein. Suitable regulatory sequences may be derived from a variety of sources, including bacterial, fungal, or viral genes (For example, see the regulatory sequences described in Goeddel, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1990). Selection of appropriate regulatory sequences is dependent on the host cell chosen, and may be readily accomplished by one of ordinary skill in the art. Examples of such regulatory sequences include: a transcriptional promoter and enhancer or RNA polymerase binding sequence, a ribosomal binding sequence, including a translation initiation signal. Additionally, depending on the host cell chosen and the vector employed, other sequences, such as an origin of replication, additional DNA restriction sites, enhancers, and sequences conferring inducibility of transcription may be incorporated into the expression vector. It will also be appreciated that the necessary regulatory sequences may be supplied by the native protein and/or its flanking regions.

The invention further provides a recombinant expression vector comprising a DNA nucleic acid molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operatively linked to a regulatory sequence in a manner which allows for expression, by transcription of the DNA molecule, of an RNA molecule which is antisense to a nucleotide sequence comprising the nucleotides as shown in Figure 7. Regulatory sequences operatively linked to the antisense nucleic

acid can be chosen which direct the continuous expression of the antisense RNA molecule.

The recombinant expression vectors of the invention may also contain a selectable marker gene which facilitates the selection of host cells transformed or transfected with a recombinant molecule of the invention. Examples of selectable marker genes are genes encoding a protein such as G418 and hygromycin which confer resistance to certain drugs, b-galactosidase, chloramphenicol acetyltransferase, or firefly luciferase. Transcription of the selectable marker gene is monitored by changes in the concentration of the selectable marker protein such as b-galactosidase, chloramphenicol acetyltransferase, or firefly luciferase. If the selectable marker gene encodes a protein conferring antibiotic resistance such as neomycin resistance transformant cells can be selected with G418. Cells that have incorporated the selectable marker gene will survive, while the other cells die. This makes it possible to visualize and assay for expression of recombinant expression vectors of the invention and in particular to determine the effect of a mutation on expression and phenotype. It will be appreciated that selectable markers can be introduced on a separate vector from the nucleic acid of interest.

The recombinant expression vectors may also contain genes which encode a fusion moiety which provides increased expression of the recombinant protein; increased solubility of the recombinant protein; and aid in the purification of a target recombinant protein by acting as a ligand in affinity purification. For example, a proteolytic cleavage site may be added to the target recombinant protein to allow separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein.

Recombinant expression vectors can be introduced into host cells to produce a transformant host cell. The term "transformant host cell" is intended to include prokaryotic and eukaryotic cells which have been transformed or transfected with a recombinant expression vector of the invention. The terms "transformed with", "transfected with", "transformation" and "transfection" are intended to encompass introduction of nucleic acid (e.g. a vector) into a cell by one of many possible techniques known in the art.

Prokaryotic cells can be transformed with nucleic acid by, for example, electroporation or calcium-chloride mediated transformation. Nucleic acid can be introduced into mammalian cells via conventional techniques such as calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofectin, electroporation or microinjection. Suitable methods for transforming and transfecting host cells can be found in Sambrook et al. (Molecular Cloning: A Laboratory Manual, 2nd Edition, Cold Spring Harbor Laboratory press (1989)), and other laboratory textbooks.

Suitable host cells include a wide variety of prokaryotic and eukaryotic host cells. For example, the proteins of the invention may be expressed in bacterial cells such as *E. coli*, insect cells (using baculovirus), yeast cells or mammalian cells. Other suitable host cells can be found in Goeddel, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1991).

The proteins of the invention may also be prepared by chemical synthesis using techniques well known in the chemistry of proteins such as solid phase synthesis (Merrifield, 1964, J. Am. Chem. Assoc. 85:2149-2154) or synthesis in homogenous solution (Houbenweyl, 1987, Methods of Organic Chemistry, ed. E. Wansch, Vol. 15 I and II, Thieme, Stuttgart).

The following non-limiting examples are illustrative of the present invention:

## EXAMPLES

### Example 1

This example demonstrates the increased expression of certain genes following pv immunization.

#### Mice:

C3H/HEJ and C57BL/6 mice were purchased from The Jackson Laboratory, Bar Harbor, ME. Mice were housed five/cage and allowed food and water *ad libitum*. All mice were used at 8-12 weeks of age.

#### Monoclonal antibodies:

The following monoclonal antibodies (Mabs) from Pharmingen (San Diego, CA) were used: anti-IL-2 (JES6-1A12; biotinylated JES6-5H4);

anti-IL-4 (11B11; biotinylated BVD6-24G2); anti-IFN $\gamma$  (R4-6A2; biotinylated XMG1.2); anti-IL-10 (JES5-2A5; biotinylated SXC-1, Pharmingen, San Diego, CA); mouse IgG1 isotype control (clone 107.3, BALB/c anti-TNP). Streptavidin horse radish peroxidase and recombinant mouse GM-CSF was also  
5 purchased from Pharmingen (San Diego, CA).

NLDC-145 (anti-mouse dendritic cells), and F(ab')<sub>2</sub> rabbit anti-rat IgG FITC conjugate (non-cross reactive with mouse IgG), or F(ab')<sub>2</sub> rabbit anti-mouse IgG PE was obtained from Serotec, Canada.

Rabbit complement, L3T4, anti-thy1.2, anti-Ly2.2, anti-Ly2.1  
10 (mouse IgG3), FITC-MAC-1 and mouse IgG1 anti-rat OX-2 were obtained from Cedarlane Labs, Hornby, Ontario.

Anti-CD28 (PV-1) and anti-CTLA (UC10-4F10-11) were obtained from Drs. C. June and J. Bluestone respectively, while anti-B7-1, anti-B7-2 were obtained from Dr. G. Powers. High titres of all 4 of the latter  
15 antibodies were produced by *in vitro* culture in a CELLMAX system (CELLCO Inc., Germantown, MD, USA).

Preparation of cells:

Spleen, Peyer's Patch (PP) and mesenteric lymph node (MLN) cell suspensions were prepared aseptically from individual mice of the  
20 different treated groups in each experiment.

Where dendritic cells were obtained by culture of bone marrow cells *in vitro* the following technique was used (Gorczynski et al., 1996a). Bone marrow plugs were aspirated from the femurs of donor male C57BL/6 (or BALB/c) mice, washed and resuspended in aF10. Cells were treated  
25 sequentially with a mixture of antibodies (L3T4, anti-thy1.2, anti-Ly2.2) and rabbit complement and dead cells removed by centrifugation over mouse lymphopaque (Cedarlane Labs, Ontario). Cells were washed x3 in aF10, and cultured in 10ml aF10 in tissue culture flasks, at a concentration of  $2 \times 10^6$ /ml with 500U/ml recombinant murine GM-CSF (Pharmingen, USA). Fresh  
30 GM-CSF was added at 36 hr intervals. Cells were separated over lymphopaque on days 3.5 and 7 of culture, again reculturing in aF10 with recombinant GM-CSF. At 10 days an aliquot of the sample was stained with NLDC-145 and FITC anti-rat IgG, anti-OX-2 and PE anti-mouse IgG,



FITC-anti-B7-1 or FITC anti-B7-2. Mean staining with these antibodies using cells harvested from such cultures has been  $93\% \pm 7\%$ ,  $14\% \pm 5\%$ ,  $78\% \pm 9\%$  and  $27\% \pm 6\%$  respectively. Remaining cells were washed, and injected into the portal vein as described.

5 Portal vein immunizations and renal transplantation:

The pv immunizations and renal transplantation were performed as described earlier (Gorczynski et al., 1994). All C3H mice received pv/iv immunization with  $15 \times 10^6$  C57BL/6 10-day cultured, bone marrow derived, dendritic cells, followed by C57BL/6 kidney transplantation. Animals received 10 intramuscular (im) injection with 10mg/Kg cyclosporin A on the day of transplantation. Mice were sacrificed for tissue harvest and RNA preparation 5 days after transplantation. In other studies animals were sacrificed as described in the text.

Where monoclonal antibodies were injected into transplanted mice, animals received 100mg intravenous (iv) at 2 day intervals (x4 injections) beginning within 2 hours of transplantation.

Cytokine production from spleen cells of transplanted mice:

In cultures used to assess induction of cytokine production spleen responder cells stimulated with irradiated (2000R) C57BL/6 spleen stimulator cells in triplicate in aF10 have been used. In multiple studies significant quantitative or qualitative differences in cytokine production from spleen, lymph node or Peyer's Patch of transplanted mice have not been seen. (Gorczynski et al., 1994b). Supernatants were pooled at 40hr from replicate wells and assayed in triplicate in ELISA assays for lymphokine production. All capture antibodies, biotinylated detection antibodies, and recombinant cytokines were obtained from Pharmingen (San Diego, CA-see above).

For IFN $\gamma$  the assay used flat-bottomed 96-well Nunc plates (Gibco, BRL) coated with 100ng/ml R4-6A2. Varying volumes of supernatant were bound in triplicate at 4°C, washed x3, and biotinylated anti-IFN $\gamma$  (XMG1.2) added. After washing, plates were incubated with streptavidin-horse radish peroxidase (Cedarlane Labs), developed with appropriate substrate and OD<sub>405</sub> determined using an ELISA plate reader.

IL-10 was assayed using a similar ELISA system with JES5-2A5 as the capture antibody, and biotinylated SXC-1 as developing antibody. Each assay reliably detected cytokine levels in the range 0.01 to 0.1 ng/ml. ELISA assays for IL-2 and IL-4 used JES6-1A12 and 11B11 as capture antibodies, with biotinylated JES6-5H4 or BVD6-24G2 as developing antibodies. Sensitivity of detection was 10pg/ml for each cytokine.

Oligonucleotide primers:

The primers used for PCR amplification for b-actin, and different cytokines, are described in previous publications (Gorczynski, R.M., 1995a; Gorczynski, R.M., 1995b; Gorczynski, R.M., 1996a). In addition, the following oligonucleotides were synthesized.

cDNA synthesis primer for driver ds cDNA (DP):  
5'-TTTTGTACAAGCTT<sub>30</sub>-3' (SEQ.ID.NO.:3)

Adapter 1 (Ad1):

5'-CTAATACGACTCACTATAGGGCTCGAGCGGCCCGCCCGGGCAGGT-3'  
(SEQ.ID.NO.:4)

Adapter 2 (Ad2):

5'-TGTAGCGTGAAGACGACAGAAAGGGCGTGGTGCGGAGGGCGGT-3'  
(SEQ.ID.NO.:5)

PCR Primer1 (P1): 5'-CTAATACGACTCACTATAGGGC-3' (SEQ.ID.NO.:6)

Nested Primer 1 (NP1): 5'-TCGAGCGGCCCGCCCGGGCAGGT-3'  
(SEQ.ID.NO.:7)

PCR Primer2 (P2): 5'-TGTAGCGTGAAGACGACAGAA-3' (SEQ.ID.NO.:8)

Nested Primer 2 (NP2): 5'-AGGGCGTGGTGCGGAGGGCGGT-3'  
(SEQ.ID.NO.:9)

Driver and Tester Preparation:

RNA was extracted from pooled mesenteric lymph node (MLN) and Peyer's Patches (PP) of 5/group renal transplant mice with iv or pv immunization. Poly(A)<sup>+</sup>mRNA was prepared from the driver (iv) group, and 2mg material used for ds cDNA synthesis with 1ng DP primer and a cDNA Synthesis Kit (Clontech) with T4 DNA polymerase. The final cDNA preparation was digested with *Rsa*I in a 50ml reaction mixture with 15 units enzyme (GIBCO) for 3 hrs, and the cDNA phenol-extracted, ethanol

precipitated, and resuspended in 7ml of deionized water (concentration approximately 300ng/ml).

*Rsa*I digested ds tester cDNA (pv group) was prepared in a similar fashion. 50ng of tester cDNA diluted in TE buffer was ligated with 2ml of Ad1 and Ad2 (each at 10mM) in separate ligation reactions at 16°C for 18hrs with 50 Units/ml T4 ligase. Thereafter 1ml of 0.2M EDTA was added, the mixture heated at 70°C for 5 min to inactivate the ligase, and the product stored at -70°C.

Subtractive hybridization and PCR amplification:

600ng driver (iv) ds cDNA was added to each of two tubes containing 20ng Ad1- and Ad2- ligated pv cDNA. The samples were mixed, precipitated with ethanol, resuspended in hybridization buffer, overlaid with mineral oil and denatured/annealed in standard fashion. The two independent samples were then combined, 200ng fresh driver cDNA added to allow for further enrichment of differentially expressed mRNAs, and the mixture again denatured and annealed for 10hrs at 68°C. The final sample was diluted in Hepes buffer with EDTA and stored at -20°C.

After subtraction two PCR amplifications were performed on the subtracted cDNA. In the first 1ml of subtracted cDNA was amplified using 1ml each of P1 and P2. The conditions for amplification were as described by Diatchenko. The amplified products were diluted 10-fold in deionized water and 1ml of product used for further amplification using the nested primers (NP1 and NP2) and a 10-cycle amplification reaction. Aliquots of the original driver/tester and subtracted cDNAs were used for PCR reactions with control oligonucleotide primers (b-actin) for known "housekeeping genes", and with primers for genes whose expression has been previously documented to be different in iv/pv immunized mice. These data are shown in Figures 1 and 2.

Figure 1 shows PCR validation of suppressive subtractive hybridization. Samples from unsubtracted (lanes 1, 3, 5 and 7) or subtracted (lanes 2, 4, 6 and 8) mRNA were reverse transcribed and tested in PCR with b-actin primers for different PCR cycle times. Lanes 1 and 2: 15 cycles; lanes 3 and 4: 20 cycles; lanes 5 and 6: 25 cycles; lanes 7 and 8: 30 cycles.

Figure 2 shows PCR validation of suppressive subtractive hybridization. Samples from unsubtracted (lanes 2 and 4) or subtracted (lanes 3 and 5) mRNA were tested as in Figure 1, except primers used were for IL-10, and different cycle times are shown. Lanes 2 and 3: 20 cycles; lanes 4 and 5: 30 cycles, lane 1: mol. wt. standard.

In addition, cloning of the subtracted cDNA was performed as follows.

Cloning and further analysis of subtracted cDNA:

The PCR amplified cDNA was cloned with a TA cloning kit (Invitrogen, California) by directly ligating into the PCR II vector. Ligation was performed at an insert:vector ratio of 3:1 in 1x ligation buffer with T4 ligase (3U/ml) overnight at 14°C. Ligation products were then inserted into INFaF' competent *Escherichia Coli* using a standard transformation protocol, and selected with ampicillin on plates containing X-gal (5-bromo-4-chloro-3-indolyl--D-galactoside). Miniprep plasmid DNA was purified with a Plasmid extraction Spin kit (Qiagen, Germany) and cut with EcoR I restriction enzyme to determine whether the plasmids contained the expected insert. Plasmids with inserts were sequenced by the dideoxy sequencing method using a T7 sequencing kit (Pharmacia Biotech, Canada). Nucleic acid homology searches were performed using the BLAST program at the National Center for Biotechnology Information (NIH, Bethesda, USA).

Further analyses of cloned material, using Northern hybridization, was as follows. Inserts in pCRII were amplified for 12 cycles using nested PCR primers. The amplified material was purified using Qiaquick Spin PCR Purification Kits (Qiagen), <sup>32</sup>P-labeled by random priming, and used as a probe for Northern hybridization with 20mg samples of the original (and fresh) iv or pv total RNA. Hybridization was performed in 5ml of ExpressHyb solution (Clontech) with a minimum of 5x10<sup>6</sup>cpm per 100ng cDNA probe and 0.1mg/ml sonicated heat-denatured salmon sperm DNA. Filters were washed 4 times, each at 15 min at 27°C with 1xSSC and 0.1% SDS, followed by a high stringency wash at 42°C for 30 min with 0.2xSSC and 0.1% SDS. Exposure times varied from 18hrs to 6 days. Figure 3 shows an autoradiograph using <sup>32</sup>P-labeled probes prepared from 4 clones obtained

using the subtraction hybridization approach described above (with pv cDNA as tester material and iv cDNA as driver). A labeled control probe was prepared with a PCR amplicon for mouse b-actin. Total RNA was prepared from mice receiving iv or pv immunization and equivalent amounts loaded in replicate lanes as shown, with gels developed from 18 hours (clone #28) to 6 days (#71). Clone 8 is most homologous with mouse poly (A) binding protein. Clone 16 is most homologous with rat MRC OX-2. Clone 28 is most homologous with human zinc-finger protein. Clone 71 has no homologous sequence.

Western blotting protocol:

The technique used was essentially that described by Sandhu et al. (1991) as modified by Bronstein et al. (1992). Samples were obtained 14 days post renal transplantation, using groups described in Figure 5. Fresh rat thymus cells were used as control. Samples were electrophoresed in 12%SDS-PAGE and transferred to PVDF membranes (Novex Co., San Diego, CA) prior to addition of primary antibody. A commercial anti-rat OX-2 was used as test reagent; control antibody was an antibody to mouse CD8a. The developing antibody used was a commercial horse-radish peroxidase labeled anti-mouse IgG. All reagents were obtained from Cedarlane Labs (Hornby, Ontario, Canada).

DNA sequence homology comparison:

Comparison of mouse OX-2 with known cDNA sequences for B7-1, B7-2, CD28 and CTLA-4 was performed using a DNASIS program (version 2.0).

**RESULTS**

Evaluation of suppression subtraction hybridization(SSH) technique

In order to evaluate the efficacy of the SSH technique used, the inventor used his previous evidence that, by PCR analysis, increased expression of mRNA for IL-10 genes was evident in lymphoid tissue from pv immunized mice. Accordingly, a dilution analysis of cDNA from the tester, driver and subtracted material, using PCR primers for b-actin and IL-10 was performed. As shown in Figure 1, after SSH there was a detectable signal for b-actin in subtracted material only after 35 cycles of amplification. By

contrast, a signal was present in the unsubtracted material after only 15 cycles. Using additional quantitative measures of template, it was found to correspond to some 1000-10,000 depletion of b-actin mRNA. In a separate study, analyzing IL-10 mRNA (Figure 2), significant enrichment of IL-10 mRNA was found as determined by comparison of the amplification detected at 30 cycles in subtracted/unsubtracted material (see lanes 4 and 5, Figure 2).

In a further test of the efficiency of subtraction the mixture of unsubtracted and subtracted tester (pv) cDNA was labeled and hybridized to Northern blots of iv (tester) and pv (driver) total RNA. The results (data not shown) indicated that the subtracted tester cDNA probe did indeed produce a significantly stronger signal with the tester RNA. Given the evidence that for any cDNA species to produce a signal in a Northern blot it must represent a concentration greater than 0.1-0.3% of the cDNA mixture, these data are again consistent with our having produced a high level of enrichment of pv-specific cDNA, with a concomitant reduction in abundant cDNAs common between tester (pv) and driver (iv) material.

Detection of unique cDNA fragments in tissue from pv immunized mice

The efficiency and validity of SSH for detection of cDNAs unique to the tissue sample from the pv immunized mice was further confirmed after cloning and sequence analysis of selected tester-specific cDNAs. 10 randomly selected cDNA clones (of 66 sequenced) were used to probe multiple preparations of pv or iv whole RNA. All revealed unique mRNAs expressed preferentially in the pv samples. Autoradiograms from 4 of these Northern blots, along with a b-actin probe as control, are shown in Figure 3. Exposure times from 18 hrs to 6 days were used which were interpreted as indicative of pv specific cDNAs of different abundance in the samples of interest.

The cDNA inserts of the 4 clones shown, along with the other 62 clones, were partially sequenced and analyzed for homology in the GenBank and EMBL data bases. A summary of these data are shown in Table 1. Note that some 30 cDNA fragments had at least 50% homology (BLAST score >250 over at least 50 nt) with other described sequences. A further 14 clones showed similar homology with known rat/human genes. Both sets may

represent members of different gene families. An additional 22 clones demonstrated no significant matches with entries in the database, and thus may represent novel genes up-regulated after pv immunization. That the data shown are a minimal estimate of such differentially expressed genes is evident from the fact that homology with IL-4 or IL-10 gene sequences (mRNAs known to be over-expressed following pv immunization-see also Figure 2) were *NOT* detected in any of the 66 clones analyzed.

The sequence homology for the clones shown in Figure 3 (>80% homology over the compared sequence) led to the further characterization of these clones. Clone 8 was shown to be most homologous with mouse poly (A) binding protein; clone 16 was shown to be most homologous with rat MRC OX-2; and clone 28 was shown to be most homologous with human zinc-finger protein. No homologous sequence was found for clone 71. In the data that follows, the analysis of one of these clones which showed homology to a rat cDNA (for OX-2, a molecule previously characterized as being preferentially expressed on rat thymocytes and dendritic cells) is described. The rationale for further investigation of this clone lies in data showing that infusion of dendritic cells via the portal vein is a potent method for prolonging allograft survival in our model systems. Note, however, that while the bone marrow derived dendritic cells that were infused via the portal vein themselves express OX-2 (see above), identical data has been obtained in Northern gels to those shown in Figure 3 using tissue harvested from mice receiving, as the earliest studies(1-5) irradiated spleen cells (OX-2<sup>+</sup> by FACS analysis) via the portal vein. In addition, in both situations, OX-2 mRNA was not detected by this suppression subtraction hybridization approach when we used tissue harvested at 0.5-2.5 days post transplantation. These results are consistent with the idea that the OX-2 signal detected is a result of novel increased expression in cells following pv immunization.

*Probing a cDNA library from tissue from pv immunized mice for expression of the murine equivalent of rat OX-2*

A cDNA library was constructed from mRNA prepared from a pool of 5 C3H mice receiving pv immunization with  $25 \times 10^6$  irradiated

(2000Rads) C57BL/6 bone marrow cells followed by renal transplantation as described in the Materials and Methods, using a kit purchased from ClonTech. Clones were plated in LB medium and probed with the <sup>32</sup>P-labeled amplicon described in Figure 3 as showing homology with rat OX-2. A 1.3Kb clone was detected, amplified, and shown after <sup>32</sup>P labeling to detect a differentially expressed product by Northern gel analysis. After sequencing using an automated DNA sequencer and fluorescent-labeled deoxynucleotides, this 1.3Kb fragment was found to share >95% homology with the region encoding the 3'untranslated region of the rat OX-2 mRNA as determined from the GeneBank sequence for rat OX-2.

Using a primer construct program, a 5'PCR primer representing positions 1-19 of the rat GeneBank sequence (corresponding to a portion of the 5'untranslated region, and the leader sequence) and 3' primers from our characterized mouse sequence were synthesized, and long-distance amplification performed to produce an amplicon predicted to encode the open-reading-frame (ORF) of the murine equivalent of the rat OX-2 gene. This amplicon was determined (as expected) to be of some 1.4Kb length. Automated sequencing produced a full-length sequence for the mouse homologue of the rat MRC OX-2 gene, including an ORF with >90% homology (predicted amino acid sequence) with the corresponding rat product, along with the 3'untranslated region. This sequence has been submitted to the Genbank (accession number AF004023).

Using a DNASIS program the predicted mouse protein sequence has some 51% homology with B7-1 and B7-2, 48% with CD28 and 54% with CTLA4 (unpublished).

*Evidence for an important role for the expressed OX-2 homologue in prolonged graft survival following pv immunization*

In an attempt to define the potential importance of the product encoded by the OX-2 gene we used a commercial antibody to rat OX-2 in a transplant model in mice receiving pv immunization and renal transplantation. In the first such study, it was asked whether there was evidence for specifically increased expression of the OX-2 molecule following pv immunization. By FACS analysis, using dual staining of hepatic mononuclear



cells and spleen cells with OX-2 and NLDC145, similar numbers of NLDC145<sup>+</sup> cells in liver or spleen samples from iv and pv immunized mice were found, ( $5 \times 10^5$  and  $6.5 \times 10^6$  respectively), but a 4-fold increase in the numbers of OX-2<sup>+</sup> NLDC145<sup>+</sup> following pv immunization. Figure 4 shows a flow cytometry profile of spleen adherent cells from iv immunized/grafted mice (panels A and B) or pv immunized/grafted mice (panels C and D). Cells were harvested 7 days after transplantation and stained with NLDC145 and F(ab')<sub>2</sub>FITC-anti-rat IgG, as well as with control (clone 107.3) mouse IgG1 serum (left hand panels) or anti-OX-2 (right hand panels) and F(ab')<sub>2</sub>PE-anti-mouse IgG. Data are representative of one of three different studies. Values shown represent the total cell population in each quadrant. The absolute numbers ( $\times 10^5$ ) of double positive cells in the liver or spleen of pv immunized mice were  $3.2 \pm 0.5$  and  $39 \pm 8$  respectively (see Figure 4 for FACS profiles of spleen adherent cells). This 4-fold increase was seen regardless of the cells used for pv immunization, either bone marrow derived dendritic cells (some 20% OX-2<sup>+</sup>-see above) or irradiated whole spleen lymphoid cells (OX-2<sup>+</sup>), suggesting that they were not merely detecting surviving OX-2<sup>+</sup> (donor) cells, but novel expression of OX-2 *in vivo*.

Western blot, Figure 5, shows increased expression of OX-2 antigen after pv immunization. The technique used for Western blotting is previously described. Samples were obtained 14 days post renal transplantation, using the groups described in Figure 6. Fresh rat thymus cells (lane 5) were used as control. Lanes 1 and 2 represent samples pooled from 3 donors/group (iv immunized; pv immunized +infusion of anti-OX-2 respectively). Samples in lanes 3 and 4 are from individual mice receiving pv immunization and renal transplantation only (no antibody treatment). Staining with anti-rat MRC OX-2 is shown in Figure 5B; with a control antibody (to mouse Ly2.1), anti-mouse cD8a, shown in Figure 5A. The developing antibody used was a commercial horse-radish peroxidase labeled anti-mouse IgG. No signal was seen using the mouse IgG1 isotype control clone 107.3 (BALB/c anti-TNP)-data not shown. Data are representative of 1 of 3 equivalent studies.

Western blotting (see Figures 5A and 5B) of samples prepared from the spleen of iv vs pv immunized and grafted mice 14 days following renal transplantation revealed staining of a band migrating with estimated molecular weight 43Kd, in agreement with data elsewhere reporting extensive glycosylation of this molecule in isolates from rat thymus. In mice receiving pv immunization along with *in vivo* treatment with anti-OX-2, no detectable signal was seen in Western blots (see lane 2, Figure 5). No staining was seen with a murine IgG1 isotype control (BALB/c anti-TNP, clone 107.3: unpublished), making it unlikely that the band observed was Fc receptor.

Figure 6 is a graft showing percent survival versus days post renal transplantation. Commercial anti-OX-2 monoclonal antibody, but not anti-mouse CD28 or anti-mouse CTLA4, reverses the graft prolongation following donor-specific pv immunization. Groups of 6 C3H mice received C57BL/6 renal allografts with no other treatment (cyclosporin A only,  $\diamond$ -), or additional pv immunization with  $15 \times 10^6$  C57BL/6 bone marrow derived dendritic cells ( $\leftarrow$ -) as described previously. Subsets of these latter mice received iv injection (every second day x4 injections) with 100mg/mouse of a commercial anti-rat OX-2 monoclonal antibody ( $\rightarrow$ -) or the isotype control (clone 107.3,  $\rightarrow$ -), or of antibodies to mouse CD28 ( $\rightarrow$ -\*) or CTLA4 ( $\rightarrow$ -\*). The animal survival for the different groups shown are pooled from 2 studies. Note that the mouse isotype control itself produced no modification of the increased renal graft survival following pv immunization. \*  $p < 0.02$ , Mann-Whitney U-test).

In two final studies mice received pv immunization and transplantation as before, but now also received iv injection with commercial anti-rat OX-2 (x4 injections; 100mg/mouse at 2 day intervals). As shown in Figures 5A and B and 6 these infusions of anti-OX-2 significantly decreased the prolonged graft survival (Figure 6) and increased expression of OX-2 antigen (Western blotting-Figures 5A and 5B) seen following pv immunization. No perturbation of graft survival following pv immunization was seen using additional treatments with anti-CD28/anti-CTLA4 (see Figure 6), or, in studies not shown, using anti-B7-1 or anti-B7-2. Again infusion of the IgG1 isotype

control Mab (clone 107.3) did not alter the increased graft survival seen following pv immunization (see Figure 6).

In separate experiments cells were harvested from mice receiving pv immunization along with additional treatment with monoclonal antibodies as show (see Table 2). Following treatment with anti-OX-2 there was no longer the altered cytokine production (with polarization to production of IL-4 and IL-10) which the inventor has described in multiple model systems in which animals received pv donor-specific pre-transplant immunization. Treatment with any of the other 4 monoclonal antibodies tested did not produce this reversal in polarization of cytokine production seen following pv immunization-indeed, using these Mabs alone in the absence of pv immunization produced a trend to increased graft survival (not shown) and significant polarization in cytokine production to increased IL-4 and IL-10 production, akin to that produced by pv immunization itself (upper half of Table 2).

OX-2 is a molecule previously characterized by Barclay et al. (1981, 1982) as being preferentially expressed on rat thymocytes and dendritic cells. Dendritic cells are known to be important signalling cells for lymphocytes, which also potentially regulate cytokine production and graft rejection, and infusion of dendritic cells is a potent means of inducing pv tolerance. The inventor has determined that OX-2 expression increased following pv immunization, and further studied whether this had any functional consequences. As shown in Figures 4 and 5, there is indeed significantly increased expression of OX-2 in spleen cells isolated from pv immunized mice, along with the increased graft survival and polarization in cytokine production (Figure 6 and Table 2). In contrast, *in vivo* infusion of anti-OX-2 abolishes increased expression of this molecule, simultaneously reversing the increased graft survival and altered cytokine profile seen. This data is consistent with the possible function of OX-2<sup>+</sup> cells in promoting allograft survival.

In the studies described the donor dendritic cells infused via the portal vein were themselves OX-2<sup>+</sup> (see description of materials and methods above). However, identical data in FACS analysis (Figure 4) and Western

Blots (Figure 5), and from suppression subtraction hybridization (Figure 3), have been obtained in studies in which we used irradiated whole spleen cells (OX-2<sup>-</sup> by FACS) for pv infusion. This is consistent with the lack of evidence for increased mRNA expression of OX-2 early (1-2 days) post transplant, as noted above. Thus it seems most likely that an operationally important “OX-2 signal” detected in the spleen of the pv immunized mice can derive from new expression, rather than necessarily from infused OX-2<sup>+</sup> cells. In the absence of a polymorphic marker for OX-2, however, it cannot be determined whether increased expression is from donor or host cells (or both). Indeed, it is perhaps somewhat surprising that the murine antibody to rat OX-2 cross-reacts in the fashion shown with murine OX-2. Definitive analysis of the *in vivo* role of OX-2 awaits similar studies to those above, using antibodies developed against the murine OX-2 homologue-these experiments are currently in progress. It is also important to point out that while pv immunization led to only a 4-fold alteration in the absolute number of detectable OX-2<sup>+</sup> NLDC<sup>+</sup> cells in the spleen/liver (see text and Figure 4), nevertheless in the face of this 4-fold difference a clear difference in OX-2 signals in Northern gels using RNA from pv vs iv immunized mice (Figure 3), along with evidence for a role for this quantitative difference in the outcome of graft survival (Figure 6) were detected. Presumably these results reflect respectively the limitation to the sensitivity of the Northern assay used, and some function of the quantitation of “co-stimulation” occurring after OX-2:OX-2 ligand interaction.

While there was some 50% homology of the predicted protein sequence of murine OX-2 with murine B7-1, B7-2, CD28 and CTLA4 (Borriello et al., 1997), antibodies to the latter molecules did not reverse the prolonged graft survival and altered cytokine production following pv immunization (Figure 6, Table 2- see also (Castle et al., 1993)). In fact these latter antibodies themselves, infused in the absence of pv immunization, produced some of the same changes in cytokine production induced by pv immunization (Table 2).

## **EXAMPLE 2**

### **Murine OX-2**

This example describes the cloning and sequencing of murine MRC OX-2.

A cDNA library was constructed from MLN cells derived from adult C3H mice, preimmunized 5 days earlier with  $10 \times 10^6$  allogeneic B10.BR bone marrow-derived dendritic cells allogeneic cells by the portal venous (pv) route, using a Cap Finder PCR cDNA library construction kit (Clontech). The inventor had previously isolated, using a PCR-Select cDNA subtraction hybridization kit (Clontech) and RNAs obtained from pooled MLN of mice immunized by the pv route or via the lateral tail vein (iv), a 350bp amplicon which showed over 98% homology with the 3' untranslated region of rat MRC OX-2 cDNA. Northern blot analysis confirmed that this amplicon detected a differentially expressed product in RNAs prepared from iv vs pv immunized mice. This amplicon was used to screen  $5 \times 10^5$  clones of the amplified library. The sequences of cDNA clones were established with an Applied Biosystems 377 Automated Sequencer, utilizing the Dye Terminator Cycle Sequencing method (Applied Biosystems, Foster City, CA). The nucleotide sequence reported in this paper has been submitted to the GenBank/EMBL Data Bank with accession number AF004023.

The cDNA shown in Figure 7 has an open reading frame of 837 base pairs, and a deduced amino acid sequence (Figure 8) of 248 amino acids, of which 30 represent a cleaved leader sequence. The predicted molecular weight of this, and the equivalent molecules in rat and human, is approximately 25kDa. The measured molecular weight in rat thymocytes, where the molecule is highly glycosylated, is 47kDa.

The murine MRC OX-2 shows some 92% and 77% homology overall at the amino acid level with equivalent molecules in rat or human respectively. As noted for the rat molecule, the sequence from a 203-229 seems likely to represent a membrane spanning domain (highly hydrophobic region), while the region from 229-248 is likely the intracytoplasmic region, with a stretch of highly basic residues immediately C-terminal to position 229. Homology in the combined transmembrane and C-terminal regions with rat and human shows some 98% and 85% similarity respectively. As predicted from membership in the Ig supergene family, there are a number of conserved Cys residues forming the disulphide bonds between b-strands of

Ig-like domains, (21 and 91; 130 and 184 respectively); residue 91 was previously found to be the most highly conserved among members of the immunoglobulin superfamily. Homology between the N-terminal Ig-domain with rat and human, versus the next Ig-domain, is 88% and 82%, or 97% and 73% respectively. This relative concentration in variability between rat and mouse in the V-terminal Ig-domain may be more understandable when the ligand specificity for the molecules in these species is clarified. Note that the presumed extracellular portion of the molecule (1-202) contains a number of sites for N-glycosylation which are preserved across species (44, 65, 73, 80, 94, 127, 130 and 151). This was previously reported for the rat cDNA sequence, and inferred from the measured size of the expressed material in rat thymocytes.

The intracytoplasmic region of the molecules has no sequence identity with known signaling kinases, nor does it have the well-described consensus sequence for the immunoreceptor tyrosine activation motif (ITAM: DXXYXXLXXXXXXXXYDXL). In addition, it lacks typical SH2 or SH3 domains to serve as "docking sites" for adapter molecules which might in turn co-opt other protein kinases in an activation cascade. Accordingly the ligand-binding activity of the extracellular domains presumably represent the biologically important region of the molecule. Some possible functions attributable to ligand interaction with OX-2 can be inferred from other data in the literature. A homologous molecule, Ng-CAM, has been reported to bind a protein-tyrosine phosphatase via N-linked oligosaccharide residues, and protein tyrosine phosphatases are known to play a key regulatory role in immune responses. More recently ALCAM, another adhesion molecule member of the Ig superfamily, the gene for which is located close to that for OX-2 on chromosome 3 in humans, has been shown to bind CD6 (a member of the scavenger receptor cystein rich family, SRCR), and antibodies to CD6 may themselves play a role in regulating immune function.

### **EXAMPLE 3**

#### **OX-2 Positive Cells Inhibit Type-1 Cytokine Production**

The inventor has shown that hepatic mononuclear, non-parenchymal, cells (NPC) can inhibit the immune response seen when allogeneic C57BL/6 dendritic cells (DC) are incubated with C3H spleen responder cells. Cells derived from these cultures transfer increased survival of C57BL/6 renal allografts in C3H mice. The inventor also found that increased expression of OX-2 on dendritic cells was associated with inhibition of cytokine production and renal allograft rejection. The inventor further explored whether inhibition by hepatic NPC was a function of OX-2 expression by these cells.

Fresh C57BL/6 spleen derived DC were cultured with C3H spleen responder cells and other putative co-regulatory cells. The latter were derived from fresh C3H or C57BL/6 liver NPC, or from C3H or C57BL/6 mice treated for 10 days by intravenous infusion of human *Flt3* ligand (*Flt3L*). Different populations of murine bone-marrow derived dendritic cells from cultures of bone marrow with (IL-4+GM-CSF) were also used as a source of putative regulator cells. Supernatants of all stimulated cultures were examined for functional expression of different cytokines (IL-2, IL-4, IFN $\gamma$ , TGF $\beta$ ). It was found that fresh C57BL/6 splenic DC induced IL-2 not IL-4 production. Cells from the sources indicated inhibited IL-2 and IFN $\gamma$  production, and promoted IL-4 and TGF $\beta$  production. Inhibition was associated with increased expression of OX-2 on these cells, as defined by semi-quantitative PCR and FACS analysis. By size fractionation, cells expressing OX-2 were a subpopulation of NLDC145+ cells. This data implies a role for cells expressing OX-2 in the regulation of induction of cytokine production by conventional allostimulatory DC.

#### **MATERIALS AND METHODS**

**Mice:** Male and female C3H/HEJ and B10.BR (H-2<sup>k/k</sup>), B10.D2 (H-2<sup>d/d</sup>) and C57BL/6 (H-2<sup>b/b</sup>) mice were purchased from the Jackson laboratories, Bar Harbour, Maine. Mice were housed 5/cage and allowed food and water ad libitum. All mice were used at 8-12 weeks of age.

**Monoclonal antibodies:** The following monoclonal antibodies (Mabs), all obtained from Pharmingen (San Diego, CA, USA) unless stated otherwise, were used: anti-IL-2 (JES6-1A12; biotinylated, JES6-5H4 ); anti-IL-4 (11B11, ATCC; biotinylated, BVD6-24G2 ); anti-IFN $\gamma$  (R4-6A2, ATCC; biotinylated XMG1.2); anti-IL-10 (JES5-2A5; biotinylated SXC-1); PE anti-B7-1/B7-2 (Cedarlane Labs, Hornby, Ontario, Canada).

Rat anti-mouse OX-2 monoclonal antibodies were prepared by Immuno-Precise Antibodies Ltd. (Victoria, BC, Canada) following immunization of rats with a crude membrane extract of LPS stimulated murine DC, followed by fusion with a non-secreting rat myeloma parent cell line (YB2/3H1.P2.G11.16Ag.20). Hybridoma supernatants were screened in ELISA using plates pre-coated with a 40-45Kd preparation of DC extracts run on Western gels (Barclay, A.N. 1981. *Immunology* 44:727; Barclay, A.N., and H.A. Ward. 1982. *Eur. J. Biochem.* 129:447). Positive clones were re-screened using FACS analysis of CHO cells transduced with a cDNA clone encoding full-length murine OX-2 (Chen, Z., H. Zeng, and R. M. Gorczynski. 1997. *BBA. Mol. Basis Dis.* 1362:6-10). FITC-conjugated F(ab')<sub>2</sub> rabbit anti-rat IgG (non cross-reactive with mouse IgG) from Serotec, Canada was used as second antibody. The Mab selected for further analysis (M3B5) was grown in bulk in a CELLMAX system (Cellco Inc., Germantown, MD). A crude preparation of rat immunoglobulin (30% saturated ammonium sulphate preparation) was used as a control Ig.

In tissue culture assays where anti-cytokine Mabs were used to confirm the specificity of the assay used 10 mg/ml of the relevant Mabs was found to neutralize up to 5.0 ng/ml of the cytokine tested.

NLDC145 (anti-mouse DC) was also obtained from Serotec. Recombinant mouse IL-4 was a kind gift from Dr. L. Yang (The Toronto Hospital); mouse rGM-CSF was purchased from Pharmingen. Recombinant human Flt3L (derived from CHO cells) was a kind gift from Dr. A.B. Troutt, Immunex Corp., Seattle, Washington, USA.



### **Renal Transplantation**

Renal transplantation was performed essentially as described elsewhere (Gorczynski, R.M. et al. 1994a. *Transplantation* 58:816-820). Animals were anesthetized with a combination of halothane and nitrous oxide inhalation, using novogesic for post-op analgesia. Orthotopic renal transplantation was performed using routine procedures. In brief, Donor animals received 200 Units of heparin, and kidneys were flushed with 2ml of ice cold heparinized physiological saline solution, prior to removal and transplantation into recipient animals with left nephrectomy. The graft renal artery was anastomosed to the recipient's abdominal aorta, and the renal artery was anastomosed to the recipient's inferior vena cava. The ureter was sewn into the recipient bladder using a small donor bladder patch. All recipients received im injection with cefotetan (30mg/Kg) on the day of transplantation and for 2 succeeding days. The remaining host kidney was removed 2 days after transplantation, unless otherwise indicated. Treatment of recipients with pv immunization, by monoclonal antibodies, or by oral immunization was as described in individual studies.

### **Portal Vein and Oral Immunization**

Portal vein and oral immunization was performed as described earlier (Gorczynski, R. M. 1995a. *Cell. Immunol.* 160:224-231; Gorczynski, R.M. et al. *Transplantation* 62:1592-1600). All animals were anaesthetized with nembutal. A midline abdominal incision was made and the viscera exposed. Cells were injected in 0.1ml through a superior mesenteric vein using a 30 gauge needle. After injection the needle was rapidly withdrawn and hemostasis secured without hematoma formation by gentle pressure using a 2mm 3 gel-foam.

Bone-marrow derived dendritic cells (DC) for pv immunization were obtained by culture of T depleted bone marrow cells *in vitro* with rIL-4 and rGM-CSF (Gorczynski, R.M. et al. *Transplantation* 62:1592-1600). Staining with NLDC145 and FITC anti-rat IgG, or with FITC anti-CD3 confirmed >95% NLDC145+ and <5% CD3+ cells at day 10 of culture (Gorczynski, R.M. et al. *Transplantation* 62:1592-1600). These cells were washed and injected into mice or used for mixed leucocyte cultures.

### Preparation of cells:

Spleen and bone marrow (Gorczynski, R.M. et al. *Transplantation* 62:1592-1600) cell suspensions were prepared aseptically from individual mice in each experiment. Hepatic mononuclear nonparenchymal cells (NPC) were isolated essentially as described elsewhere (Gorczynski, R.M. 1994b. *Immunology* 81:27-35). Tissue was first digested at 37°C for 45 min with a mixture of collagenase/dispase, prior to separation (15 min at 17,000rpm at room temperature) over mouse lymphopaque (Cedarlane Labs). Mononuclear cells were resuspended in a-Minimal Essential Medium supplemented with 2-mercaptoethanol and 10% fetal calf serum (aF10). Where cells were obtained from Flt3L injected mice, animals were treated by iv injection of 10mg/mouse Flt3L daily for 10 days. After enzyme digestion recovery of liver/spleen cells from these mice was markedly increased compared with saline-injected controls ( $120 \times 10^6$ ,  $390 \times 10^6$  vs  $7 \times 10^6$  and  $120 \times 10^6$  respectively).

### Cytotoxicity and cytokine assays:

In cultures used to assess induction of cytotoxicity or cytokine production responder cells were stimulated with irradiated (2000R) stimulator cells in triplicate in aF10. Supernatants were pooled from replicate wells at 40 hrs for cytokine assays (below). No reproducible differences in cytokine levels have been detected from cultures assayed between 36 and 54 hrs of stimulation. In some experiments the cultures received 1mCi/well (at 72 hrs) of  $^3\text{HTdR}$  and proliferation was assessed by harvesting cells 14 hrs later and counting in a well-type b-counter.

Where cytotoxicity was measured cells were harvested and pooled from equivalent cultures at 5 days, counted, and recultured at different effector:target with  $^{51}\text{Cr}$  EL4 ( $\text{H2}^{\text{b/b}}$ ) or P815 ( $\text{H2}^{\text{d/d}}$ ) tumor target cells. Supernatants were sampled at 4 hrs for assessment of specific cytotoxicity.

IL-2 and IL-4 activity were assayed by bioassay using the IL-2/IL-4 dependent cell lines, CTLL-2 and CT4.S respectively. Recombinant cytokines for standardization of assays was purchased from Genzyme (Cambridge, MA). IL-2 assays were set up in the presence of 11B11 to block potential stimulation of CTLL-2 with IL-4; IL-4 assays were set up in the

presence of S4B6 to block IL-2 mediated stimulation. Both the IL-2 and IL-4 assays reproducibly detected 50 pg of recombinant lymphokine added to cultures.

In addition, IL-2, IL-4, IFN $\gamma$  and IL-10 were assayed using ELISA assays. For IFN $\gamma$  the assay used flat-bottomed Nunc plates (Gibco, BRL) coated with 100ng/ml R4-6A2. Varying dilutions of supernatant were bound in triplicate at 4°C, washed x3, and biotinylated anti-IFN $\gamma$  (XMG1.2) added. After washing, plates were incubated with streptavidin-horse radish peroxidase (Cedarlane Labs, Hornby, Ontario), developed with appropriate substrate, and OD<sub>405</sub> determined using an ELISA plate reader. Recombinant IFN $\gamma$  for standardization was from Pharmingen. IL-10 was similarly assayed by ELISA, using JES5-2A5 as a capture antibody and biotinylated SXC-1 as developing antibody. rIL-10 for standardization was from Pepro Tech Inc. (Rocky Hill, NJ). Each assay detected 0.1ng/ml cytokine. ELISA assays for IL-2 and IL-4 used JES6-1A12 and 11B11 as capture antibodies, with JAS6-5H4 or BVD6-24G2 as developing antibodies. Sensitivity of detection was 20pg/ml for each cytokine. Where checked the correlation between bioassay and ELISA for IL-2 or IL-4 was excellent ( $r>0.90$ ). In all studies reported below, data are shown from ELISA assays only. Where cytokine data are pooled from several studies (e.g. Figures 14, 16, 17), absolute values of cytokine production were obtained as above using commercial recombinant cytokines to standardize the assays. In our hands, supernatants from C3Hanti-C57BL/6 cultures, under the conditions described, reproducibly contain  $950\pm200$  and  $80\pm25$  pg/ml IL-2 and IL-4 respectively.

#### **Preparation of RNA:**

Different sources of tissue from renal-grafted female mice receiving DC and kidney allografts from male mice were harvested for RNA extraction as described elsewhere (Gorczynski, R. M. 1995a. *Cell. Immunol.* 160:224-231). The OD<sub>280/260</sub> of each sample was measured and reverse transcription performed using oligo (dT) primers (27-7858: Pharmacia, USA). The cDNA was diluted to a total volume of 100ml with water and frozen at -70°C until use in PCR reactions with primers for murine GAPDH, B7-1, B7-2 or OX-2. The sense (S) and antisense (AS) primers were synthesized by the

Biotechnology Service Centre, Hospital for Sick Children, Toronto, using published sequences. 5' primers were <sup>32</sup>P end-labeled for PCR and had comparable levels of specific activity after purification by ethanol precipitation. 5ml cDNA was amplified for 35 cycles by PCR, and samples were analyzed in 12.5% polyacrylamide gels followed by overnight (18 hrs) exposure for autoradiography. In control studies, using H-Y primer sets, this technique reliably detects H-Y mRNA from extracts of female spleen cells to which male cells are added at a concentration of 1:105 (Gorczynski, R. M. 1995a. *Cell. Immunol.* 160:224-231; Gorczynski, R.M. et al. *Transplantation* 62:1592-1600). Quantitative comparison of expression of different PCR products used densitometric scanning of the autoradiograms.

GAPDH Sense: 5'TGATGACATCAAGAAGGTGGTGAAG3' (SEQ.ID.NO.:10)

GAPDH Antisense: 5'TCCTTGGAGGCCATGTAGGCCAT3' (SEQ.ID.NO.:11)

B7-1 Sense: 5'CCTTGCCGTTACAACCTCTCC3' (SEQ.ID.NO.:12)

B7-1Antisense: 5'CGGAAGCAAAGCAGGTAATC3' (SEQ.ID.NO.:13)

B7-2 Sense: 5'TCTCAGATGCTGTTTCCGTG3' (SEQ.ID.NO.:14)

B7-2 Antisense: 5'GGTTCACTGAAGTTGGCGAT3' (SEQ.ID.NO.:15)

OX-2 Sense: 5'GTGGAAGTGGTGACCCAGGA3' (SEQ.ID.NO.:16)

OX-2 Antisense: 5'ATAGAGAGTAAGGCAAGCTG3' (SEQ.ID.NO.:17)

## **Statistical analysis:**

In studies with multiple groups, ANOVA was performed to compare significance. In some cases (as defined in individual circumstances) pairwise comparison between groups was also subsequently performed.

## **RESULTS**

### **Antigen stimulation, in the presence of hepatic NPC, induces development of a cell population capable of inhibiting proliferation and IL-2 production on adoptive transfer:**

In a previous manuscript (Gorczynski, R. M. et al., *Transplantation*. 66: 339-349) it was reported that C3H spleen cells stimulated in the presence of syngeneic NPC and allogeneic (C57BL/6) DC produced a cell population able to inhibit generation of IL-2 from fresh spleen cells stimulated with C57BL/6 DC, and capable of inhibiting C57BL/6 renal allograft

rejection *in vivo*. In order to ask whether this function of NPC was MHC restricted or not, the following study was performed.

5 C57BL/6 (H2<sup>b/b</sup>) spleen cells were stimulated *in vitro* with B10.BR (H2<sup>k/k</sup>) bone-marrow derived DC, in the presence/absence of the following NPC: C57BL/6; B10.BR; B10.D2 (H2<sup>d/d</sup>). In addition, control cultures were incubated with the NPC only. Proliferation and IL-2/IL-4 production was measured in one aliquot of these primary cultures. In addition, at 5 days, cells were harvested from another set of the primary cultures, washed, and  $2 \times 10^5$  cells added to cultures containing  $5 \times 10^6$  fresh  
10 C57BL/6 spleen cells and B10.BR DC. Proliferation and cytokine production was measured in these latter cultures in standard fashion. Data pooled from three equivalent studies are shown in panels A) and B) of Figure 9.

Figure 9 is a bar graph showing regulation of proliferation and cytokine production following stimulation by allogeneic DC using hepatic NPC in accordance with the methods described herein. In panel A) cultures were  
15 initiated with  $5 \times 10^6$  C57BL/6 responder spleen cells alone (group 1), or with  $2 \times 10^5$  B10.BR DC (group 2). Further groups (3-5, and 6-8 respectively) contained C57BL/6 responder cells and  $2 \times 10^5$  NPC from either C57BL/6, B10.D2 or B10.BR respectively (3-5) or these same NPC and B10.BR DC (6-8). Data show mean proliferation and cytokine production from triplicate  
20 cultures in three separate studies. In panel B) data show proliferation and cytokine production from cultures of  $5 \times 10^6$  C57BL/6 responder spleen cells stimulated in triplicate with  $2 \times 10^5$  B10.BR DC alone, or with the addition also of  $2 \times 10^5$  cells harvested from the cultures shown in the upper panel. Again  
25 data represent arithmetic means of 3 separate experiments. \*  $p < 0.05$  compared with control cultures (far left in each panel).

There are a number of points of interest. As previously documented, addition of NPC syngeneic with spleen responder cells (C57BL/6 in this case) to cells stimulated with allogeneic (B10.BR) DC led to  
30 decreased proliferation and IL-2 production from those responder cells compared with cells stimulated by DC alone (compare groups 6 and 2 of upper panel of Figure 9, panel A). IL-4 production in contrast was enhanced. NPC alone, whether syngeneic or allogeneic to the responder cells, produced

no obvious effect (groups 3-5, panel A) of Figure 9). Furthermore, cells from primary cultures receiving the DC+NPC mixture were able to inhibit proliferation and IL-2 production (while promoting IL-4 production) from fresh spleen cells stimulated in secondary cultures with the same (B10.BR) DC (see panel B) of Figure 9). However, data in this Figure make another important point. The same inhibition of proliferation/IL-2 production in primary cultures was seen using either B10.BR NPC (MHC matched with the DC stimulus-group 8, panel A) of Figure 9) or with third-party B10.D2 NPC (MHC-mismatched with both spleen responder cells and allogeneic stimulator DC-group 7, panel A) of Figure 9). Again no obvious effect was seen in cultures stimulated with B10.BR or B10.D2 NPC alone (groups 4 and 5). Finally, cells taken from primary cultures stimulated with DC and NPC from either B10.BR or B10.D2 could also inhibit proliferation/IL-2 production from secondary C57BL/6 spleen cell cultures stimulated with B10.BR DC-again cells taken from primary cultures with NPC alone produced no such inhibition (see panel B) of Figure 9). Thus the inhibition of proliferation/IL-2 production and enhancement of IL-4 production seen in primary cultures, as well as the induction of suppression measured in secondary cultures, all induced by NPC, are not MHC-restricted.

**Specificity of inhibition/suppression induced by hepatic NPC:**

One interpretation of the data shown in Figure 9 and elsewhere is that NPC deliver a signal to DC-stimulated cells which is distinct from the antigen-signal provided by the DC themselves (and is MHC non-restricted). This signal modulates the antigen-specific signal provided by the DC. In order to assess the antigen-specificity of the immunoregulation described in Figure 9, the following experiment was performed.

C57BL/6 spleen responder cells were stimulated with B10.D2 or B10.BR bone marrow-derived DC, in the presence/absence of NPC from B10.BR or B10.D2 mice. Proliferation and cytokine production was measured in aliquots of these cultures as before. In addition, further aliquots of cells harvested from these primary cultures were added to cultures of fresh C57BL/6 spleen cells stimulated with B10.BR (panel B)-Figure 10) or B10.D2

(panel C)-Figure 10) DC. Again proliferation and cytokine production was measured. Data pooled from three such studies are shown in Figure 10.

Figure 10 shows specificity of inhibition of proliferation of cytokine production by hepatic NPC (see Figure 9 and description of Figure 9 for more details). In panel A),  $5 \times 10^6$  C57BL/6 spleen cells were stimulated in triplicate for 3 days with  $2 \times 10^5$  B10.BR or B10.D2 DC, with/without  $2 \times 10^5$  NPC derived from B10.D2 or B10.BR mice. Data shown are arithmetic means of 3 repeat studies. In panels B) and C), fresh C57BL/6 responder spleen cells were cultured in triplicate with either B10.BR DC (panel B), or B10.D2 DC (Panel C), with/without  $2 \times 10^5$  additional cells from the primary cultures (groups 1-6 in panel A). Again data represent arithmetic means of proliferation/cytokine production from 3 studies. \*  $p < 0.05$  compared with control cultures (far left in each panel).

Data from the primary cultures (panel A)) recapitulates the observations made in Figure 9, and show that NPC inhibit proliferation and IL-2 production from DC-stimulated responder cells in an antigen and MHC-unrestricted fashion. However, the data in panels B) and C) of this figure show clearly that adoptive transfer of inhibition using cells from these primary cultures occurs in an antigen-restricted fashion, dictated by the antigen-specificity of the DC used in the primary cultures, not of the NPC used for induction of suppression. These auxiliary cells in the NPC population thus have a functional property of being "facilitator cells for induction of suppression". Note that in other studies (data not shown) where the final assay system involved measuring cytotoxicity to allogeneic target cells, a similar inhibition of lysis (rather than cytokine production) was seen using cells harvested from primary cultures stimulated with DC and hepatic NPC (see Gorczynski, R. M., et al. 1998a. *Transplantation*. 66: 000-008).

**Hepatic cell preparations from Flt3L treated mice are a potent source of DC and "facilitator" cells:**

It has been reported at length that pv infusion of alloantigen, or iv infusion of liver-derived allogeneic mononuclear cells induces operational unresponsiveness in recipient animals (Gorczynski, R. M. 1995a. *Cell. Immunol.* 160:224-231; Gorczynski, R.M. et al. *Transplantation* 62:1592-

1600; Gorczynski, R.M. et al. 1994a. *Transplantation* 58:816-820.;  
Gorczynski, R. M., and D. Wojcik. 1992. *Immunol. Lett.* 34:177-182;  
Gorczynski, R. M. et al. 1995b. *Transplantation*. 60:1337-1341). The total  
hepatic mononuclear cell yield from normal mice is of the order of  $5 \times 10^6$   
5 cells/mouse. In order to increase the yield, and explore the possibility that the  
liver itself might be a source both of allostimulatory DC and "facilitator" cells 2  
C57BL/6 mice were exposed for 10 days to daily iv infusions of 10mg/mouse  
human CHO-derived *Flt3L*, a known growth factor for DC (Steptoe, R. J. et al.  
1997. *J Immunol.* 159:5483-5491). Liver tissue was harvested and pooled  
10 from these donors and mononuclear cells prepared as described in the  
Materials and Methods section above (mean  $130 \times 10^6$  cells/donor). These  
cells were further subjected to sub-fractionation by size using unit gravity  
sedimentation techniques (Miller, R. G., and R. A. Phillips. 1969. *J. Cell.*  
*Comp. Physiol.* 73:191-198). A typical size profile for recovered cells is  
15 shown in Figure 11 (one of 3 studies).

Figure 11 shows OX-2 expression in a subpopulation of NPC. It  
is a sedimentation analysis (cell profile) and FACS analysis of cells isolated at  
10 days from *Flt3L*-treated C57BL/6 mice. Two C57BL/67 mice received  
10mg/mouse *Flt3L* iv daily for 10 days. Hepatic NPC were sedimented for 3  
20 hrs at 4°C, and the fractions shown collected (Fxs 1-4 with sedimentation  
velocities 2.5-3.8, 3.8-5.1, 5.1-6.4 and 6.4-8.0 mm/hr respectively). Aliquots  
of the cells were stained in triplicate with the Mabs shown. The remainder of  
the cells were used as in Figures 12-14. Data are pooled from 3 studies.

In these same studies cells isolated from the various fractions  
25 shown in Figure 11 were tested as follows. Firstly, cells were stained with  
FITC-labeled Mabs to B7-1, B7-2, NLDC145 and rat anti-mouse OX-2 (M3B5)  
with FITC anti-rat IgG as second antibody. In addition, mRNA extracted from  
the different cell samples were assayed by PCR for expression of GAPDH,  
B7-1, B7-2 and OX-2. Data are shown in Figures 11 (pooled from 3 separate  
30 studies) and Figure 12 (representative PCR data from one experiment).

Figure 12 shows PCR detection of B7-1, B7-2 and OX-2 in  
hepatic NPMC. It is a PCR analysis for mRNA expression of OX-2, B7-1 and



B7-2 in various hepatic NPC cell fractions isolated from *Flt3L* treated mice (see Figure 11). Data are representative from 1 of 3 studies.

Further aliquots of the cells were used to stimulate fresh C3H spleen responder cells in culture. Proliferation and cytokine assays were performed as before (see Figure 9), and in addition cells were taken from these primary cultures and added to fresh secondary cultures of C3H spleen responder cells and C57BL/6 bone marrow-derived DC. Again proliferation and cytokine production was assayed from these secondary cultures. Data pooled from 3 studies of this type are shown in Figure 13 (panels A) and B).

Figure 13 shows that hepatic NPMC from *Flt3L* treated mice results IL-2 and IL-4 production. Stimulation of proliferation/cytokine production by NPC from *Flt3L* treated mice, and inhibition of the same (where stimulation is induced by a separate population of DC) is a function of different cell populations. (See text and Figures 11-12 for more details.) Hepatic NPC fractions were derived from *Flt3L* treated C57BL/6 mice and were used to stimulate C3H spleen cells in triplicate cultures, alone or in the presence of bone-marrow derived C57BL/6 DC (see panel A). Data show arithmetic means for proliferation/cytokine production from 3 experiments. In addition, cells harvested from these primary cultures were added to fresh C3H spleen cells stimulated with C57BL/6 DC (panel B), and again proliferation/cytokine production assayed. \*  $p < 0.05$  compared with control groups (far left of panel).

Finally, cells from the various fractions were infused iv into 2/group C3H mice which also received C57BL/6 renal allografts as antigen challenge. Spleen cells were harvested from these individual mice 10 days after transplantation and restimulated in culture with C57BL/6 or B10.D2 DC, again with cytokines measured at 40 hrs (see Figure 14).

Figure 14 is a bar graph of cytokines produced from cells from C3H mice with C57BL/b renal allografts and NPC from *Flt3* treated C57BL/6 donors. OX-2<sup>+</sup> NPC infused iv into renal transplant allograft recipients leads to polarization of cytokine production (to IL-4, IL-10 and TGF $\beta$ ) in spleen cells harvested from those mice and restimulated *in vitro*. Fractions of NPC from *Flt3L* treated C57BL/6 mice (from Figure 11) were infused iv into 2/group C3H

recipients, receiving C57BL/6 renal allografts (along with CsA) in standard fashion (see Materials and Methods). Mice were sacrificed 14 days after transplantation and spleen cells stimulated *in vitro* in triplicate with C57BL/6 DC stimulator cells. Cytokines were assayed in the supernatants of these cultures at 60hrs. Data show arithmetic means pooled from cultures in 3 studies of this type. \*  $p < 0.05$  compared with control groups (far left-no NPC infused).

Data in Figure 11 show that distinct subpopulations of slow-sedimenting cells express OX-2 in the cells harvested from Flt3L treated mice, when compared with cells expressing B7-1 and/or B7-2. In general expression of OX-2 and B7-2 occurred in equivalent subpopulations. Faster-sedimenting cells (Fx 3 and 4 in Figure 11), while staining for NLDC145, were positive by fluorescence mainly for B7-1, not B7-2 or OX-2. Similar conclusions were reached both by FACS analysis of cell populations (Figure 11) , and by PCR analysis of mRNA (Figure 12).

When the functional capacity of these different cell populations was investigated (Figures 13 and 14) it was found that optimal direct stimulation (or proliferation and IL-2 production) was seen from B7-1 expressing cells (Fxs 3 and 4 in panel A) of Figure 13), while only OX-2 expressing cells (Fxs 1 and 2 in Figures 11 and 12) were capable of producing the inhibitory effects defined earlier (Figures 9 & 10) in the two-stage culture system (panel B) in Figure 13). These same cells (Fxs 1 and 2) were in turn able, after iv infusion, to polarize cells from mice given renal allografts to produce predominantly IL-4, IL-10 and TGF $\beta$  production on restimulation *in vitro* (Figure 14). These data are consistent with the notion that after FltL treatment of mice expansion of a population of immunostimulatory DC occurs within the liver, which also contains another distinct population of (facilitator) cells which promote immunoregulation.

**Evidence that cell populations with “facilitator” activity from the liver of Flt3L treated mice prolong graft survival *in vivo*:**

Since it has been reported elsewhere that there is a good correlation between treatments (such as pv immunization) which decrease IL-2 production and increase IL-4 production from restimulated cells and

prolongation of graft survival (Gorczynski, R. M., and D. Wojcik. 1994. *J. Immunol.* 152:2011-2019; Gorczynski, R. M. 1995a. *Cell. Immunol.* 160:224-231; Gorczynski, R.M. et al. *Transplantation* 62:1592-1600), and that increased expression of OX-2 is also independently associated with increased graft survival after pv immunization (Gorczynski, R. M. et al. 1998b. *Transplantation.* 65:1106-1114), the next question was whether cells isolated from Flt3L treated mice which induced inhibitory function *in vitro* (see Figures 9, 10 and 13), and expressed increased amounts of OX-2 (Figures 11, 12) were themselves capable of promoting increased graft survival *in vivo*.

Groups of 2 C57BL/6 mice received iv infusions of 10mg/mouse Flt3L for 10 days as before. Cells were isolated from the liver by enzyme digestion, and fractionated by unit gravity sedimentation. 4 pools of cells were recovered, and an aliquot stained as before in FACS with anti-OX-2. Groups of 2 C3H mice received  $10 \times 10^6$  cells iv from the 4 separate pools. A control group received saline injections only. Over the next 48 hrs all mice received C57BL/6 renal transplants. All mice received CsA (10mg/Kg) on the day of renal transplantation. Data in Figure 15 are pooled from 3 studies of this type (representing 6 mice/group), and show the animal survival in these 5 different groups.

Figure 15 shows NPC from Flt3L treated C57BL/6 mice, infused iv into recipient C3H mice, inhibit C57BL/6 renal allograft rejection. Two mice groups received the different subpopulations of NPC derived from Flt3L treated mice shown in Figures 11 and 12. Fxs 1 and 2 were OX-2+. Mice received C57BL/6 renal allografts within 48 hrs along with CsA (see Materials and Methods). Animal survival was followed as an end point. Data shown are pooled from 3 studies (6 mice/group). \* $p < 0.05$  compared with mice receiving CsA only ().

It is quite clear from this Figure that only hepatic cells expressing OX-2 (Fxs 1 and 2-see Figures 11 and 12) were capable of promoting increased graft survival after iv infusion. Comparison of these data with those in Figure 13 confirm that these cell populations were also those identified, using a 2-stage culture assay system, as cells with functional "facilitator" activity (see also Figures 9 and 10). There was no significant

difference in survival between groups receiving NPC-Fx1 or NPC-Fx2 in this experiment, in keeping with relatively equivalent levels of OX-2 expression in these fractions (Figure 11).

**Anti-OX-2 monoclonal antibody *in vitro* reverses regulation induced by hepatic NPC:**

A final study was directed to whether anti-OX-2 monoclonal antibody M3B5, added to cultures of C3H spleen responder cells, allogeneic (C57BL/6) DC and NPC from C57BL/6 mice, could prevent the inhibition of IL-2 production in primary cultures, and the development of cells able to inhibit such cytokine responses from freshly stimulated responder cells in secondary cultures (see Figures 9, 10 and 13). Data in Figures 16 and 17 are pooled from 3 studies of this type.

Figure 16 is a bar graph showing the effect of anti B7-1; B7-2; or OX-2 on primary allostimulation. It shows that anti-OX-2 Mab increases IL-2 cytokine production *in vitro* after stimulation of C3H responder spleen cells with C57BL/6 DC. Subgroups of cultures contained the Mabs shown. Cytokines were assayed at 60 hrs. All data represent arithmetic means pooled from 3 repeat studies. \* $p < 0.05$  compared with control group (far left).

Figure 17 is a bar graph showing that anti-OX-2 reverses inhibition by NPC. It shows that anti-OX-2 Mab inhibits development of immunoregulatory cells *in vitro* following incubation with hepatic NPC. C3H responder spleen cells were incubated in triplicate with C57BL/6 DC along with NPC (see Figures 9 and 10). Subgroups of these cultures contained the Mabs shown. Cytokines were assayed in cultures at 60 hrs (panel A). In addition, cells were harvested from all groups, washed and added to fresh C3H responder spleen cells and C57BL/6 DC (panel B). Cytokines in these groups were assayed 60 hrs later. All data represent arithmetic means pooled from 3 repeat studies. \* $p < 0.05$  compared with control group from cultures of NPC with no monoclonal antibodies (far left in Figure)-see also Figure 16.

Primary cultures were of two types, containing C3H responder spleen cells and C57BL/6 DC alone (Figure 16), or the same mixture with added C57BL/6 NPC (Figure 17). Subsets of these cultures contained in

addition either 5mg/ml of anti-B7-1, anti-B7-2 or anti-OX-2. Supernatants from responder cells stimulated in the presence of DC only were assayed after 60hrs for cytokine production (Figure 16). For the primary cultures incubated with both DC and NPC, supernatants were harvested at 60hrs and tested for cytokine production (Figure 17A). In addition, cells were harvested after 5 days, washed, and added to secondary cultures of fresh C3H responder cells with fresh C57BL/6 DC. No monoclonal antibodies were added at this second culture stage. Data for cytokine production these secondary cultures are shown in Figure 17B.

Addition of anti-B7-1 or anti-B7-2 to DC stimulated spleen cultures led to inhibition of cytokine production (Figure 16), while in contrast anti-OX-2 monoclonal antibody led an increase in IL-2 production in these primary cultures (Figure 16). We have reported similar findings elsewhere (Ragheb et al-submitted for publication). Interestingly, anti-OX-2 abolished the inhibition of cytokine production caused by NPC in these primary cultures (Figure 17A-see also Figures 9, 10 and 13). In addition, anti-OX-2 prevented the functional development of a cell population capable of transferring inhibition of cytokine production to freshly stimulated spleen cells (Figure 17B).

## **DISCUSSION**

There is considerable theoretical as well as practical interest in understanding the mechanism(s) by which a state of antigen specific tolerance can be induced in lymphoid populations. Limits to the effective induction of tolerance represent a major challenge to more successful allo (and xeno) transplantation, to name but one example (Akatsuka, Y., C. Cervený, and J. A. Hansen. 1996. *Hum. Immunol.* 48:125-134). Significant efforts have been invested into exploring how pre- (or peri-) transplant donor-specific immunization might produce such a state (Qian, J. H. et al. 1985. *J. Immunol.* 134:3656-3663; Kenick, S., et al. 1987. *Transpl. Proc.* 19:478-480; Gorczynski, R. M. 1992. *Immunol. Lett.* 33:67-77; Thelen, M., and U. Wirthmueller. 1994. *Curr. Opin. Immunol.* 6:106-112; Akolkar, P. N. et al. 1993. *J. Immunol.* 150 (April 1):2761-2773; Ahvazi, B. C. et al. *J. Leu. Biol.* 58 (1):23-31; Albina, J. E. et al. 1991. *J. Immunol.* 147:144-152). There

is good evidence that portal venous (pv) immunization somehow leads to tolerance induction, and this immunoregulation can apparently be monitored by following changes in cytokine production from host cells, with decreased production of IL-2, IL-12 and IFN $\gamma$ , and increased IL-4, IL-10, IL-13 and TGF $\beta$  ( Thelen, M., and U. Wirthmueller. 1994. *Curr. Opin. Immunol.* 6:106-112; Gorczynski, R. M. et al. 1998a. *Transplantation.* 66: 000-008). Which, if any, of these cytokine changes is directly and causally implicated nevertheless remains obscure.

Further analysis of the cell population able to induce tolerance after pv immunization led to the somewhat paradoxical observation that donor dendritic (DC) cells represented an excellent tolerizing population (Gorczynski, R. M. 1995a. *Cell. Immunol.* 160:224-231; Gorczynski, R.M. et al. *Transplantation* 62:1592-1600). Since antigen-pulsed DC are conventionally thought of as representing an optimal immunizing regime, the mechanism(s) activated following DC pv immunization which led to tolerance (Banchereau, J., and R. M. Steinman. 1998. *Nature.* 392:245-252) was of interest. It is already clear that DC themselves represent an extremely heterogeneous population, in terms of origin, cell surface phenotype, turnover *in vivo* and possibly function (Salomon, B. et al. 1998. *J. Immunol.* 160:708-717; Leenen, P. J. M. et al. 1998. *J. Immunol.* 160:2166-2173). In the mouse lymph node at least 3 discrete populations were identified, one of which comprised small CD8a<sup>+</sup>NLDC145<sup>+</sup> cells, likely of lymphoid origin, with an immature phenotype, and whose numbers were profoundly increased (100x) following *Flt3L* treatment *in vivo* (Salomon, B. et al. 1998. *J. Immunol.* 160:708-717) (administration of the latter has been reported to lead to proliferation of dendritic cells and other cells of hematopoietic origin (Maraskovsky, E. et al. 1996. *J. Exptl. Med.* 184:1953-1962)). These cells resembled the interdigitating DC found in the T cell areas of the splenic white pulp, and have been implicated in regulation of immunity induced by other (myeloid derived) DC (Salomon, B. et al. 1998. *J. Immunol.* 160:708-717; Kronin, V. et al. 1996. *J. Immunol.* 157:3819-3827; Suss, G., and K. Shortman. 1996. *J. Exptl. Med.* 183:1789-1796).

A variety of other studies have indicated that the induction of immunity (vs tolerance) following antigen presentation was intrinsically dependent upon the co-existence of other signaling ligands at the surface of DC (interacting with appropriate counter-ligands on the surface of other cells (e.g. stimulated T cells)) (Larsen, C. P. et al. 1994. *J. Immunol.* 152:5208-5219; Lenschow, D. J. et al. 1996. *Annu. Rev. Immunol.* 14:233-258; Larsen, C. P., and T. C. Pearson. 1997. *Curr. Opin. Immunol.* 9:641-647). It was speculated that infusion of DC via the portal vein induced tolerance by co-opting another cell population, distinguishable by expression of unique cell surface ligands, whose biological function was to facilitate induction of tolerance, not immunity, when antigen was presented in association with otherwise immunogenic DC. Some preliminary evidence supporting this hypothesis was recently reported (Gorczynski, R. M. et al. 1998a. *Transplantation.* 66: 339-349). Herein, this is referred to as a facilitator cell. Moreover, because pv immunization has been shown to be associated with increased expression of a novel molecule, OX-2, previously reported to be expressed on DC (Barclay, A.N. 1981. *Immunology* 44:727; Barclay, A.N., and H.A. Ward. 1982. *Eur. J. Biochem.* 129:447; Chen, Z. et al. 1997. *BBA. Mol. Basis Dis.* 1362:6-10; Gorczynski, R. M. et al. 1998b. *Transplantation.* 65:1106-1114), it was predicted that this molecule would in fact serve as a "marker" for the hypothetical facilitator cell described. Experiments reported herein are consistent with such a hypothesis.

It is here shown that within the hepatic NPC population there is a subset of cells able to inhibit stimulation by allogeneic DC in a non-MHC restricted fashion (see Figures 9 and 10), and able to induce the development of an antigen-specific immunoregulatory cell population *in vitro* (see Figures 9 and 10). The non-MHC-restricted nature of this "facilitator" cell interaction indicates that it functions by providing an accessory signal (a regulatory not a co-stimulatory signal) to the DC which stimulate T cells in the allogeneic mixed leukocyte reaction described, in a fashion analogous to the original description of costimulatory interactions (Jenkins, M. K. et al. 1988. *J. Immunol.* 140:3324-3329). As a result the stimulated lymphocytes alter their cytokine production profile (with decreased IL-2 production and proliferation),

and become able to regulate the immune response seen from freshly stimulated lymphocytes (see panel B in Figures 9 and 10). Most interestingly, following expansion of DC *in vivo* by *Flt3L* treatment, it is shown that in fact the liver itself contains both an immunostimulating population (large cells by velocity sedimentation analysis), and this putative “facilitator” cell population (see Figures 11-15). Furthermore, the latter biological activity resides within a slow-sedimenting (small size) NLDC145<sup>+</sup> cell population expressing preferentially both cell surface B7-2 and OX-2 (see Figures 11 and 12). When it was investigated whether this same population of cells was active *in vivo* in regulating graft tolerance, it was found again that after prior *Flt3L* treatment, the liver contained a population of cells which transferred increased renal graft acceptance (Figure 15) and in parallel altered the cytokine production profile of immunized mice towards increased IL-4 and TGF $\beta$ , and decreased IL-2 and IFN $\gamma$  production (Figure 14).

In a final attempt to explore the role for OX-2 expression itself in this regulatory function, fresh spleen cells were stimulated with DC alone or in the presence of anti-B7-1, anti-B7-2 or anti-OX-2. Note that other studies (data not shown) have confirmed that even the bone-marrow derived DC used contains small numbers of OX-2<sup>+</sup> cells (RMG-unpublished). Unlike anti-B7-1 and anti-B7-2 which decreased cytokine production, a result in keeping with the hypothesized role for these as costimulator molecules (Hancock, W. W. et al. 1996. *Proc. Natl. Acad. Sci. USA*. 93:13967-13972; Freeman, G. J. et al. 1995. *Immunity*. 2:523-532; Kuchroo, V. K. et al. 1995. *Cell*. 80:707-718), anti-OX-2 produced a small but significant (1.7-2.5 fold in three studies) increase in IL-2 production in this system (Figure 16). Most important, however, inclusion of anti-OX-2 Mab in a system where exogenous “facilitator” cells were added (from NPC), blocked completely the induction of inhibition normally seen in such cultures (Figures 9 and 10; compare with lower panel of Figure 17). These data are consistent with the concept that OX-2 delivers a regulatory, not a costimulatory, signal in this situation.

How does the present data fit within the evolving framework of understanding in the heterogeneity of DC? As noted above, there has been speculation that a separate population of CD8a<sup>+</sup>NLDC145<sup>+</sup> DC of lymphoid



origin which proliferates in response to *Flt3L*, might be responsible for immunoregulation. Other data have implicated IL-10 as a cytokine which might modify development/maturation of DC into a population expressing increased amounts of B7-2 and capable of inducing tolerance (Steinbrink, K. et al. 1997. *J Immunol.* 159:4772-4780). The role of regulation of expression of Fas as a controlling feature in this regard is unexplored (Suss, G., and K. Shortman. 1996. *J. Exptl. Med.* 183:1789-1796). The data disclosed herein is the first to implicate another molecule, OX-2, in the delivery of a tolerizing signal, perhaps in association with alterations in expression of B7-2, Fas etc. It is intriguing that while there is clearly a key role for intra-thymic DC in the regulation of self-tolerance (Banchereau, J., and R. M. Steinman. 1998. *Nature.* 392:245-252), natural expression of OX-2 was initially first described on thymic DC (as well as within the brain) (Barclay, A.N. 1981. *Immunology* 44:727)-there is as yet no evidence to suggest that this represents a functionally relevant expression for OX-2 in this location. However, other independent data have also implied an immunoregulatory role for OX-2 expression, again as assayed by altered cytokine production *in vitro* from cells stimulated in the presence/absence of expressed OX-2 (Borriello, F. et al. 1997. *J. Immuno..* 158:4548).

It has been reported that following pv immunization there is a measureable expansion in numbers of populations of  $\gamma\delta\text{TCR}^+$  cells capable of adoptive transfer of increased graft survival to naïve recipients (Gorczynski, R. M. et al. 1996c. *Immunology.* 87 (3):381-389). Little is known concerning the nature of the antigen recognized by these cells, and why, as a population, their numbers are preferentially increased following pv immunization. It is speculated that this may be explainable ultimately in terms of a differential susceptibility of  $\gamma\delta\text{TCR}^+$  vs  $\alpha\beta\text{TCR}^+$  cells to immunoregulatory signals delivered following OX-2 expression.

In conclusion, the inventor has reported for the first time that functional heterogeneity in the DC pool may be understandable in terms of differential expression of OX-2 on the cell surface. Expression of this molecule seems to give cells the capability to induce immunoregulation, increased renal graft survival (and altered cytokine production both *in vivo*

and *in vitro*). The present invention suggests that such OX-2 expressing cells are referred to as "facilitator" cells (for tolerance induction).

#### **EXAMPLE 4**

##### **Preparation of Murine Antibodies**

5                    Mouse and rat hybridomas to a 43Kd molecule expressed in the thymus, on a subpopulation of dendritic cells, and in the brain, in mammalian tissue derived from mouse, rat and human were prepared. Using CHO cells transiently transfected with adenovirus vector(s) expressing a cDNA construct for the relevant OX-2 gene, the monoclonal antibodies (Mabs) detect a  
10                   molecule encoded by this construct (rat OX-2 (rOX-2), mouse OX-2 (mOX-2) and human OX-2 (huOX-2) respectively). Furthermore, at least some of the anti-rat Mabs detect determinants expressed on the murine OX-2 molecule.

##### **MATERIALS AND METHODS**

Antigen preparation from tissues and Western blotting were performed as  
15                   described in Gorczynski et al., Transplantation, 1998, 65:1106-1114:

                    Spleen cells (human samples were obtained from cadavers at the time of organ retrieval for transplantation) were used for preparation of dendritic cells/macrophages. Tissue was digested with a mixture of collagenase and dispase and centrifuged over lymphopaque. Cells were  
20                   adhered for 2 hr at 37°C, washed vigorously, and incubated for 14 hr at 37°C. Dendritic cells were isolated as non-adherent cells (Gorczynski et al., Transplantation, 1996. 62:1592-1600). Routine staining of mouse splenocytes with NLDC-145 and FITC anti-rat IgG, or FITC-MAC-1 before and after overnight incubation produced the following staining pattern in these  
25                   adherent cells: 8%±2%, 90%±11% and 92%±9%, 9%±3% respectively. The crude (non-adherent) dendritic cell preparation was extracted with lysis buffer, titred to a protein concentration of 10mg/ml, and used for immunization. Some of the same material was used subsequently in screening ELISAs (below).

30                   When brain tissue was used in Western gel analysis, whole tissue extract was electrophoresed in 12%SDS-PAGE and transferred to PVDF membranes (Novex Co., San Diego, CA). Putative anti-OX-2 Mabs were used as test reagent, with isotypic antibodies (negative in ELISA tests)

used as controls. Membranes were developed using either anti-rat or anti-mouse horse radish peroxidase and appropriate substrate.

Immunization and production of Mabs:

Four female BALB/c mice were initially immunized by intraperitoneal injections with 1mg of human or rat dendritic antigen in Complete Freund's Adjuvant. Three subsequent boosts were administered as above, spaced at 3 week intervals, with Incomplete Freund's Adjuvant. When the serum titre had risen more than 10-fold from a pre-immune serum sample, as determined by ELISA, the 2 highest responders were boosted intravenously. Three days later the donor mice were sacrificed and the spleen cells were harvested and pooled. Fusion of the splenocytes with X63-Ag8.6.5.3 BALB/c parental myeloma cells was performed as previously described (Kohler, G. and C. Milstein. 1975. *Nature*. 25: p. 256-259), except that one-step selection and cloning of the hybridomas was performed in 0.8% methylcellulose medium (Immuno-Precise Antibodies Ltd., Victoria, BC). This proprietary semi-solid medium allows HAT selection and cloning in a single step and eliminates the overgrowth of slower growing desirable clones by faster growing, perhaps undesirable, hybridomas. Clones were picked and resuspended in wells of 96-well tissue culture plates in 200 µl of D-MEM medium containing 1% hypoxanthine/thymidine, 20% Fetal Bovine serum, 1% OPI, and  $1 \times 10^6$ /ml BALB/c thymocytes. After 4 days, the supernatants were screened by ELISA for antibody activity on plates coated with the immunizing antigen. Putative positive hybridomas were re-cloned by limited dilution cloning to ensure monoclonality and screened in FACS on extracts prepared from brain tissue (below).

For the production of rat mAbs, 2 Fisher rats were immunized as above with mouse antigen. Essentially the same procedure was followed, except the parental cell line used for the fusion was YB2/0.

ELISA and FACS analysis of putative Mabs:

ELISA assays used polystyrene plates pre-coated with 100ng/ml poly-L-lysine, followed by overnight incubation with the crude dendritic cell antigen (used for immunization) at 10mg/ml. Wells were developed after binding of hybridoma supernatants using the anti-rat/anti-mouse horse radish

peroxidase antibodies above and plates were analysed in an automatic ELISA plate reader (TiterTek Multiskan, MCC/340, FlowLabs, Mississauga, Ontario, Canada).

FACS analysis was performed using putative anti-OX-2 Mabs and the following cells. Fresh peripheral blood leucocytes (PBL), isolated over rat/mouse lymphopaque (Cedarlane laboratories) or Ficoll-Hypaque (human); fresh spleen dendritic cells (isolated after adherence and overnight incubation, as above); and CHO cells transduced with viral vectors engineered to contain a single copy of a cDNA inserted into the not1/bamH1 sites, encoding the relevant species-specific OX-2, as per published sequences (Chen, Z. et al. 1997. *BBA. Mol. Basis Dis.* 1362:6-10; McCaughan, G.W., et al. 1987. *Immunogenetics*. 25: p. 133-135), or with control vector alone. FITC anti-mouse (or anti-rat) IgG was used as secondary antibody.

Mixed leucocyte reactivity (MLR) and cytokine production:

Allogeneic MLR cultures, using 1:1 mixtures of  $2.5 \times 10^6$  responder PBL and mitomycin C treated stimulator PBL, were set up in 24-well culture plates in 1ml of aMEM medium supplemented with 10% FCS. Cells were obtained from C3H responder mice (with stimulator C57BL/6), Lewis (LEW) rats (with Brown Norway, BN, as stimulator), and individual human donors. Culture supernatants were harvested at 60 hrs and tested for different cytokines using previously described ELISA assays (mouse), or using CTLL-2 as bioassay for IL-2 production from all responder cell sources (Gorczynski, R.M., et al. 1998c. *Immunology*. 93: p. 221-229).

**RESULTS**

Evaluation of a number of Mabs for staining of cell populations in fresh PBL or spleen:

All Mabs tested in the experiments herein described were previously screened as described in the Materials and Methods above, and detected a molecule in Western gel of brain extracts with Molecular Weight 42-45 Kd, and also stained CHO transduced by OX-2 encoding viral vectors. Data in Table 3 show FACS analysis for these Mabs using fresh cells. The data are summed over several independent analyses, using a number of

Mabs directed to rat, mouse or human OX-2, for staining of cells harvested from fresh PBL or spleen (adherent cells only were tested for the latter: these represented some 5%-8% of the total cell population in all cases).

It is clear from Table 3 that PBL in all species tested contained some 1.3%-2.5% OX-2<sup>+</sup> cells by FACS analysis, and that spleen adherent cells similarly contained 4%-8% OX-2<sup>+</sup> cells. As confirmation of the inventor's previous work, spleen adherent cells taken from C3H mice or LEW rats treated 4 days earlier by portal venous immunization with 20x10<sup>6</sup> (or 50x10<sup>6</sup> respectively) of C57BL/6 (or BN) bone marrow cells showed some 3.5-5 fold elevation in OX-2<sup>+</sup> cells (see Table 3). Under these conditions specific increases in survival of subsequent allo-transplanted cells/tissue have been reported (Gorczyński, R.M. et al. 1996a. *Transplantation* 62:1592-1600).

Ability of anti-OX-2 Mabs to modulate cytokine production in MLR *in vitro*:

In a final study the issue of whether these Mabs can modify the immune response (as assayed by cytokine production) of cells stimulated in an allogeneic mixed leucocyte reaction (MLR) *in vitro* was addressed. The inventor has previously shown that cells taken from mice pretreated by portal allogeneic immunization produce predominantly type-2 cytokines, and that an anti-OX-2 Mab could apparently reverse this polarization in cytokine production (and indeed abolish the increased graft survival seen in such mice). Data in Table 4 confirm these results using 3 independent Mabs to mouse OX-2. Further, rat or human cells stimulated in the presence of anti-rat (or human) OX-2, similarly show more pronounced IL-2 production than cells stimulated in the presence of isotypic control Ig (or no Ig), without a generalized increase in cytokine production (as analysed here by no change in IL-6 production in any group).

**DISCUSSION**

In the data in this example it is confirmed that using species specific Mabs, to human, rat or mouse OX-2, that Mabs to the molecule detected on the surface of host dendritic cells may play a role in regulating cytokine production after allostimulation *in vitro*, and more particularly that functionally blocking OX-2 expression leads to enhanced IL-2 production (a type-1 cytokine) after allostimulation (Table 4). Borriello et al also recently

reported that OX-2 expression was not a costimulator for induction of IL-2 and IFN $\gamma$  synthesis (Borriello, F. et al. 1997. *J. Immuno.* 158:4548)-our data imply it is in fact a negative signal for type-1 cytokine production. In mice preimmunized by the portal vein, as reported earlier, there is a 4-fold increase in OX-2 expressing cells in PBL and spleen, and a reversal of polarization in cytokine production (from type-2 cytokines to type-1 cytokines) after stimulation of cells in the presence of OX-2 (see Tables 3 and 4) (Gorczynski, R. M. et al. 1998b. *Transplantation*. 65:1106-1114).

### **EXAMPLE 5**

#### **Preparation of Rat Antibodies**

Five rats were immunized using GERBU adjuvant (GERBU Biotechnik, Gaiberg, Germany) with 500mg of membrane protein purified from the mouse dendritic cell (DC) line DC2.4 (a gift from K. Rock, Harvard). Serum from these rats was tested 7 days after the third immunization, and compared with a pre-immunization sample in an ELISA using plate-bound material of Mol. Wt. 40Kd-45Kd eluted from Western blots, and Alk Pase anti-rat Ig. Two rats with high titre antibody were re-immunized and sacrificed 4 days later for fusion of spleen cells with HAT-sensitive Sp2/0 parent cells for preparation of hybridomas. Hybridomas were screened by ELISA (56/960+ve), subcloned, and frozen (-70°C). For further specificity testing of the anti-OX-2 Mabs will use CHO cells can be transfected with a pBK eukaryotic expression vector (Stratagene,CA) expressing OX-2. Full length OX-2 cDNA, including the leader sequence, was amplified from DC2.4 cells using sense and antisense primers constructed with Spe1 or Xba1 sites respectively at their 5' ends for directional cloning into the vector. A band of the expected size (849bp) was obtained on agarose gel electrophoresis. The sequence of the cloned cDNA was confirmed by sequencing using an automated DNA sequencer (Chen, Z. and Gorczynski, R.M. 1997. *Biochem. Biophys. Acta*. 100, in press). CHO cells were transfected by electroporation (5x10<sup>6</sup> cells in 0.5ml were pulsed at 960MHz and 120V using a Bio-Rad Gene Pulser (Bio-Rad, Hercules, CA), using the full length OX-2 expression plasmid along with a plasmid encoding puromycin resistance (100:1 ratio), followed by selection in puromycin (12mg/ml for 4 days). Puromycin resistant cells were

cloned by limiting dilution. 5 CHO transfectant clones have been obtained expressing mRNA for OX-2 as confirmed by PCR. These clones can be used to screen the putative rat anti-mouse OX-2 Mabs.

(a) FACS staining of cells from pv immunized mice with anti-mouse OX-2

5 A 4-fold increase in staining of spleen and hepatic NLDC145+ (dendritic cell marker) cells from pv immunized mice with anti-rat OX-290 was observed. Spleen and liver tissue of mice at various times (12 hours; 2, 7 and 14 days) following pv immunization can be sectioned and stained by immunohistochemistry, using anti-NLDC145, anti-OX-2 Mabs. Single cell  
10 suspensions from the same tissues can be stained, using 3-colour FACS, with FITC-anti-mouse OX-2, rhodamine-anti-NLDC145, and phycoerythrin-anti-T200 (mouse lymphocyte marker). In all cases (both FACS and immunohistochemistry) the appropriate irrelevant isotype control antibodies are included. Tissue from control mice receiving renal grafts alone, or  
15 following additional iv immunization, can also be examined. Detection of NLDC145+ (and/or MAC-1+) cells showing increased expression of OX-2 is predicted in pv immunized mice only (see Gorczynski, R.M. et al. 1998. *J. Immunol.* 160, in press). The inventor has shown DC-associated antigen persists only in animals with surviving grafts (Gorczynski, R.M., Chen, Z., Zeng, H. and Fu, X.M. 1998. *Transplantation* submitted). It was also  
20 assessed whether anti-OX-2, infused at different times post transplantation, causes rejection (b).

(b) Modulation of graft rejection and cytokine production by anti-mouse OX-2

25 C3H mice receive pv immunization with cultured C57BL/6 bone-marrow derived dendritic cells (DC), CsA and renal allografts. Groups of mice receive intravenous infusion of various rat anti-mouse OX-2 Mabs (100-500mg/mouse, x5, at 2 day intervals), beginning at different times post transplantation (this will be guided by data from (a)). Serum creatinine and animal survival are followed. Serum from Mab-treated mice are tested in  
30 ELISA and by FACS with OX-2 expressing CHO transfectants (above) to ensure antibody excess. If OX-2 expression is important for pv induced increased graft survival, the anti-OX-2 treated pv immunized mice will reject grafts like untreated controls, with similar polarization of cytokine production

to type-1 cytokines (assayed by PCR; ELISA with cultured, restimulated cells). As controls pv immunized, grafted mice receive anti-CD28 and anti-CTLA4 these Mabs do not modify the effects of pv immunization as assayed by graft survival or polarization in cytokine production. It is expected that OX-2 treatment but not other Mabs, will simultaneously abolish expansion of  $\gamma\delta$ TCR+ cells after pv immunization.

#### **EXAMPLE 6**

##### **Preparation of a fusion protein linking the extracellular domain of OX-2 to mouse Fc**

Immunoadhesins, in which a hybrid molecule is created at the cDNA level by fusing the extracellular domain (ED) of an adhesion molecule with the carboxyl terminus of IgG heavy chain, the whole being expressed in mammalian cells or in a baculovirus system, have been powerful tools in the identification and isolation of the counter ligands for the adhesion molecule of interest. Ligands for a number of members of the TNFR family, were identified in this fashion (Goodwin, R.G. et al. 1993. *Eur. J. Immunol.* 23, 2631-2641; Gruss, H. and Dower, S. 1995. *Blood* 85, 3378-3404). Interest has developed in the potential application of immunoadhesins as therapeutic agents. A CTLA4 immunoadhesion, with the capacity to bind both B7-1 and B7-2, has been used to inhibit T cell costimulation and decrease rejection (Larsen, C.P. et al. 1996. *Nature* 381, 434-438). Note that CD28/CTLA4 are not counter ligands for OX-289. The fusion protein, is predicted to alter cytokine production (increased IL-4, IL-10; decreased IL-2, IFN $\gamma$ ) and increase renal graft survival like pv immunization. We expect that synergistic blockade of costimulation (e.g. by CTLA4-Fc) and triggering of a coregulatory pathway (by OX-2ED-Fc) will induce tolerance and produce indefinite graft survival.

##### **a) Construction of an OX-2 fusion protein with murine IgGFc2a**

A cDNA encoding the extracellular region of OX-2 (OX-2ED) was amplified by PCR, using a 5' oligonucleotide primer which inserts a Sal1 site 5' immediately at the start of the V-region sequence and a 3' primer which creates a BamH1 site at the 3' end (the site of junction with Fc). Using cDNA prepared from mouse ConA activated spleen cells, with a 5' primer containing an Spe1 site, and a 3' primer containing a Sal1 site, the signal peptide for IL-6



(SP-IL-6) was amplified by PCR and ligated to the OX-2ED amplicon. In frame ligation across the junction of SP-IL-6 and OX-2ED was checked by manual sequencing-the final cDNA amplified by the 5'SP-IL-6 primer and the 3'OX-2ED primer was, as expected, 695bp. A plasmid expressing murine IgGFC2a (Fcγ2a), modified to create a unique BamH1 site spanning the first codon of the hinge region, and with a unique Xba1 site 3' to the termination codon, has been obtained from Dr. Terry Strom (Zheng, X.X. et al. 1995. *Journal of Immunology*. 154, 5590-5600). The IgGFC2a in this insert has been further modified to replace the C1q binding motif (rendering it non-lytic) and inactivate the FcγR1 binding site (see Zheng, X.X. et al. 1995. *Journal of Immunology*. 154, 5590-5600). Ligation of OX-2ED and IgGFC2a in the correct reading frame at the BamH1 site yields a 1446bp long open reading frame encoding a single 478-amino acid polypeptide (including the 24-amino acid IL-6 signal peptide). The homodimer has a predicted 105kDa Mol Wt, exclusive of glycosylation. The fusion gene is then cloned as an Spe1-Xba1 cassette into the eukaryotic expression plasmid pBK/CMV (Stratagene, CA). This plasmid has a CMV promoter/enhancer and a neomycin-resistance gene for selection using G418. The appropriate genetic construction of the OX-2ED-Fc can be confirmed by direct sequencing after cloning into the plasmid vector (Chen, Z. and Gorczynski, R.M. 1997. *Biochem. Biophys. Acta*. 100, in press)-see also above. The plasmid is transfected into CHO cells by electroporation (see above), and selected in medium with 1.5mg/ml G418 (Geneticin:Life Technologies, Inc.). After subcloning, high producing clones are selected by screening culture supernatants in ELISA using anti-OX-2 Mabs as capture antibody, and Alk Pase coupled anti-IgGFC2a as detection antibody. OX-2ED-Fc fusion protein is purified from culture supernatants using protein A-Sepharose affinity chromatography, dialysed against PBS, filter-sterilized and stored in aliquots at -20°C. The size, and OX-2 (+IgGFC2a) specificity of the secreted product can be confirmed using Western blot analysis under reducing (+DTT) and non-reducing (-DTT) conditions, with Mabs to OX-2 and rat monoclonal anti-mouse IgGFC2a (Pharmingen). The product can be titrated as an inhibitor for FACS staining of OX-2 expressing CHO cells (see above) using rat Mabs to OX-2 as probe.

As a prelude to studies (below) using OX-2ED-Fc *in vivo*, the half-life ( $t_{1/2}$ ) in mouse serum following injection of groups of 6 8-week C3H mice will be studied. This is carried out by subjecting mice to iv injections of 50mg or 10 mg of OX-2ED-Fc, and obtains serial 50ml blood samples at 0.3, 1, 6, 24, 48, 72 and 96 hours. The serum is analyzed in ELISA using plates coated with anti-OX-2 as capture antibody, and Alk Pase coupled monoclonal anti-IgGFc2a for detection (thus ensuring the assay detects only OX-2ED-Fc, not OX-2 or IgGFc2a alone). Based on earlier data in which Fc fusion proteins were used to extend the *in vivo* half-life, a  $t_{1/2}$  in the range of 30-40 hrs (Zheng, X.X. et al. 1995. *Journal of Immunology*. 154, 5590-5600) is predicted.

b) OX-2: IgGFc Immunoadhesion Inhibits MLR

CHO cells were transduced with a vector carry the OX-2:Fc cDNA insert. Supernatant was harvested from the CHO cells at 7 days and was cultured with  $5 \times 10^6$  LEW spleen and  $2.5 \times 10^6$  irradiated LBNFI spleen cells. The supernatant contained 50ng/ml OX-2:Fc.

The results, shown in Table 5, demonstrate that the soluble OX-2:Fc immunoadhesion inhibits IL-2 production and generation of cytotoxic T cells and induces IL-4 production. These results support the use of OX-2 as an immunosuppressant.

c) Use of OX-2:Fc *in vivo* for Prevention of Graft Rejection

It was shown in (b) that incubation in the presence of 50ng/ml OX-2:Fc can inhibit an *in vitro* MLR reaction. To detect inhibition of *in vivo* graft rejection, C3H mice received C57BL/6 skin grafts along with iv injection of OX-2:Fc (50mg/mouse) every 2 days x4 injections. Grafts were inspected daily after 10 days for rejection. In a separate study 3 mice/group (receiving saline or OX-2:Fc) were sacrificed at 10 days and spleen cells restimulated *in vitro* (x48 hrs) for analysis of cytokine production. Data for these studies is shown in Tables 6 and 7. It is clear from these data that OX-2:Fc has the potential for use as an immunosuppressant to prolong graft acceptance. Furthermore, in association with increased graft survival in this model, OX-2:Fc alters polarization in cytokine production, as already described for portal vein donor-specific immunization.

## **EXAMPLE 7**

### **OX-2 Expression in Placenta**

Using *in situ* hybridization, the inventor has shown that OX-2 is not expressed in the placenta of mice with increased potential for fetal loss. In contrast, OX-2 is expressed in the placenta of normal, non-aborting mice.

CBA/J and DBA/2J mice were used. Matings of CBA/J(females) with DBA/2J males show a high incidence of fetal loss (>80%), unlike the reverse scenario. Placental tissue was obtained from matings at 8.5 days of gestation. Uteri were snap frozen, 5mm sections cut, and stained with a biotinylated anti-sense probe for murine OX-2. Data shown in Figures 18A and 18B indicate increased expression of OX-2 mRNA (in situ labeling) in the non-aborting strain combination, with essentially absent expression in the aborting combination. These data are consistent with the notion that OX-2 expression prevents spontaneous fetal loss syndrome.

The data show that there are fewer OX-2+ implantation sites on day 8.5 of pregnancy in mice which are predisposed to fetal loss syndrome (CBAxDBA/2 matings) by contrast to CBAxBALB/c matings which are not so predisposed. Fgl2 is the trigger for loss, and where OX-2 is also expressed, these potentially doomed implantations are "rescued". This follows from the finding that the abortion rate is lower than expected from % fgl2++ implantation sites, unless anti-OX-2 mAb is administered. In the latter instance, the abortion rate rises to equate with the estimated proportion of fgl2++ implant sites.

## **EXAMPLE 8**

### **OX-2 Prevents Fetal Loss**

Successful pregnancy in allopregnant mice can also be viewed as dependent upon control of graft rejection. Proinflammatory Th1 cytokines (TNF- $\alpha$  + IFN- $\gamma$  + IL-1) can cause spontaneous abortion in mice by a mechanism which involves a novel prothrombinase, fgl2, which promotes fibrin deposition. However, the inventors found that spontaneous abortion rates in abortion-prone CBA x DBA/2 matings and in low abortion rate CBA x BALB/c matings were lower than the frequency of implantation sites showing fibrin<sup>hi</sup> + fgl2 mRNA<sup>hi</sup>. OX2 expression was present in the same sites as fgl2

mRNA, and neutralization of this OX2 expression by anti-OX-2 raised the abortion rate to predicted levels. Conversely, an OX2 immunoadhesin dramatically reduced the abortion rate. Therefore, in addition to its role in organ and tissue allograft rejection, OX2 expression is involved in the prevention of spontaneous abortion triggered by cytokine up-regulation of fgl2 at the feto-maternal interface.

In the Example detailed below, evidence is shown using anti-OX2 monoclonal antibody and a soluble form of OX2 in which the extracellular domains of the molecule are linked to an Ig Fc region (OX2:Fc) (described in Example 6), that OX2 is fundamentally important to achieving successful allopregnancy.

## **METHODS**

All of the techniques used, including mixed leukocyte cultures, cytokine analysis, and allografting, are detailed in previous publications (Gorczynski et al. 1996a; Arck et al. 1997a; and Clark 1999). The anti-OX2 mAb (3B6) was obtained from BioCan (Mississauga, Ontario) (Gorczynski et al. 1998b). 100µg/mouse was used for each injection. A polyclonal, affinity-purified, rabbit antibody to fgl2 was described elsewhere, and used ip at a dose of 22mg/mouse. OX2Fc immunoadhesin was given ip (35µg/mouse).

## **RESULTS**

### ***In situ expression of OX2 mRNA following renal transplantation or allopregnancy:***

OX-2 has been reported at the fetomaternal interface using immunohistochemistry in rats (Example 7 and Bukovsky et al. 1984). To determine whether OX-2 was expressed in the uterus of allopregnant mice, the inventors carried out in situ hybridization for OX-2 mRNA in CBA x DBA/2 and CBA x BALB/c matings. Adjacent sections of the tissue samples were also used to stain for fgl2 mRNA (fgl2 is a prothrombinase molecule up-regulated by certain Th1 cytokines implicated triggering pregnancy loss). For comparison, we also examined OX2 expression in liver sections from C3H mice receiving C57BL/6 renal allografts following donor-specific portal vein preimmunization, a treatment which promotes tolerance and which is critically dependent upon up-regulation of expression of OX-2 on hepatic APC. Typical

patterns for uterine staining (pregnant mice) are reported elsewhere (Clark et al. 2001), along with cumulative data for OX2 and fgl2 expression in the uteri of pregnant control mice and mice treated with  $\text{TNF}\alpha$  +  $\text{IFN}\gamma$  to increase abortion rates (Clark et al. 2001).

5 In these studies the inventors found that OX2 mRNA expression was up-regulated following pv immunization and renal transplantation, and in  
10 allopregnant mice. In pregnant mice the inventors found a negative correlation between expression of the molecules fgl2 and OX-2 which did not reach statistical significance (Clark et al. 2001). With following cytokine treatment of pregnant animals, and prior to the onset of abortions, the proportion of fgl2<sup>hi</sup>  
15 implantations increased, although this also did not achieve significance due to small numbers. However, with cytokine treatment, the proportion of OX-2<sup>hi</sup> implants decreased dramatically. These data support the hypothesis that in pregnancy, fgl2 and OX-2 expression are reciprocally regulated by cytokines, that their levels affect pregnancy outcome, and that a major determinant of success or failure of fgl2<sup>hi</sup> 'at risk' implantations was the presence or absence of OX2. Most interestingly, the inventors also reported that continued expression of OX2, as occurs in pregnancy, was essential for successful survival of allografts following pv pretransplant immunization, and for the  
20 concomitant changes in cytokine production seen in those animals (previous Examples and Gorczynski et al. 2000).

Effect of anti-OX2 mAb on renal transplant survival and pregnancy outcome:

25 Fifty to 70% of implantations in control and cytokine-boosted CBA x DBA/2 pregnancies show the fgl2<sup>hi</sup> phenotype (Clark et al. 2001). An injection of anti-Vg1.1 on day 8.5 of pregnancy, 1 day before abortions become evident, inactivates most of the trophoblast-recognizing  $\gamma\delta$  subset producing IL-10 and  $\text{TGF}\beta$  and boosts abortion rates to approximately 48% (Arck et al. 1999). The inventors hypothesized that the "suppressor"  $\gamma\delta\text{T}$  cells inactivated by this treatment might be dependent upon OX2 expression for  
30 their functional activity. The inventors have reported that following allotransplantation, the kinetics of expression of OX2 follows closely the development of immunoregulatory  $\gamma\delta\text{T}$  cells. Functional blockade of OX2 expression (by anti-OX2 treatment) reverses increased graft survival (see

Figure 2) and prevents adoptive transfer of tolerance by  $\gamma\delta$ T cells. To test whether OX-2 expression in pregnancy might similarly be activating anti-abortion mechanisms which 'rescued' fgl2<sup>hi</sup> implant sites from proceeding to embryo death where both OX-2 and fgl-2 were expressed, the inventors injected control CBA x DBA/2- and CBA x BALB/c-mated mice with the same anti-OX-2 monoclonal antibody that blocks induction of transplantation tolerance. Figure 19 shows that injection on or after day 8.5 increased the spontaneous abortion rate to that expected if all fgl2<sup>hi</sup> sites in CBA x DBA/2 proceeded to resorb. The increase in abortion rate was not due to a toxic effect of anti-OX-2 on the embryo because co-administration of anti-fgl2 to neutralize prothrombinase activity abrogated the boost in abortion rates (data not shown). Injection of anti-OX-2 into CBA x BALB/c-mated mice also increased the abortion rate to 22 %, consistent with the 21% fgl2 mRNA<sup>hi</sup> implantation site frequency.

*Infusion of OX2 immunoadhesin modulates renal allograft rejection and spontaneous abortion:*

As further "proof of principle" that OX2 expression is functionally important for increased allograft survival the inventors used an immunoadhesin (OX2:Fc), in which the extracellular domains of OX2 were linked to a murine IgGFc region, to investigate modulation of allograft rejection and pregnancy. Data in the previous Examples has already indicated that this molecule has potent immunoregulatory properties in vivo, including the ability to decrease allograft rejection. Data in Figure 20 represent a cumulative comparison of the effect of infusion of OX-2:Fc on renal allograft survival or rate of abortions in CBA x DBA/2 mated mice. Once again there was a clear parallel between the functional activity of OX2:Fc measured in these two assays.

**DISCUSSION**

The notion that the rejection of organ allografts would be mimicked immunologically by immune recognition of the fetus in allopregnant mothers has been with us for decades. There has been intense interest in the role of altered cells and soluble factors (e.g.cytokines) in the phenomena seen in both situations. As an example, with decreased graft rejection (and

successful pregnancy) there are numerous reports of the presence of unique  $\gamma\delta$ T cells with "suppressor phenotypes", and altered cytokine patterns, with elevated levels of type-2 cytokines, in particular IL-10 and TGF $\beta$  (Chaouat et al. 1999; Tsuda et al. 2001; Clark et al. 2001b; Arck et al. 1999 and Arck et al. 1997a). In contrast, allograft rejection in rodents and man has been associated with elevated type-1 cytokines, and previous studies have also shown that both TNF $\alpha$  and IFN $\gamma$  must be present for spontaneous abortions to be induced (Chaouat et al. 1999; Tsuda et al. 2001; Clark et al. 2001b; Arck et al. 1999 and Arck et al. 1997a). Since the fgl2 gene is activated by IFN $\gamma$  but not by TNF $\alpha$ , the inventors have hypothesized that the obligatory role for the latter cytokine either involves activation of PMNL essential for abortions to be completed, or down-regulates OX-2 expression.

It is worthy of note that in the inventors' previous work, and in the studies described above, expression of fgl2, a thrombosis-inducing molecule, was up-regulated on the trophoblast in response to cytokines. Cytokine-treated IRF1<sup>-/-</sup> females mated to +/+ males do not abort, as there is no up-regulation of fgl2 in the maternal decidual tissues and the best explanation for lack of abortions is that fgl2<sup>+</sup> trophoblast and fgl2<sup>+</sup> maternal decidua must meet. In the regions where the two tissues meet, a zone of spontaneous cleavage, enough enzymatic activity presumably occurs to cause necrosis. The inventors have also documented a basal level of fgl2 expression in trophoblast tissue. Given the evidence that excessive anticoagulation with heparin or hirudin leads to retroplacental hemorrhage fatal to the embryo and sometimes mother, the inventors suggest that this low (basal) level of expression of the molecule fgl2 may reflect a normal homeostatic role for fgl2 in preventing spontaneous bleeding. However, cytokine-mediated (by TNF $\alpha$  and IFN $\gamma$ ) upregulation of fgl2 is associated with increased rates of abortion. These effects can in turn be counteracted by the combination of both TGF- $\beta$  and IL-10, both perhaps produced by trophoblast cells, which are known to inhibit cell-mediated vascular injury and clotting. There is little data to date examining the role of fgl2 prothrombinase in allograft rejection. Interestingly xenograft rejection, a process in which acute and subacute vascular changes are believed crucial, is reportedly less

pronounced in an fgl2 knockout mouse (Levy et al, personal communication). It thus becomes extremely interesting to know whether fgl2 has a more general role in immunomodulation in both allo and fetal grafts.

The present data shed further light on the mechanisms by which some of these changes occur in both the allopregnant mouse and in allo-transplant, by providing evidence for a crucial role for altered expression of another molecule OX2, in regulating both embryo execution triggered by cytokine up-regulation of fgl2 prothrombinase and the modulation of renal allograft rejection. The inventors have proposed that OX2 acts in transplantation as a co-stimulatory signal that deviates cytokine production away from Th1 (e.g. IL-2, IFN- $\gamma$ ) and towards Th2/3 (e.g. IL-4, IL-10, TGF- $\beta$ ) production. Associated with this is expansion of a  $\gamma\delta$  T cell subset that mediates tolerance via suppression. In support of such a hypothesis, the inventors have shown increased expression of OX2 is associated with decreased rejection and altered cytokine production; that the kinetics of expression of OX2 parallels altered cytokine production and  $\gamma\delta$ T cell expansion; and that these effects are diminished by infusion of anti-OX2 mAbs, and enhanced by infusion of the immunoadhesin OX2:Fc. The inventors have now also documented that a similar correlation exists between OX2 expression and fetal loss in allopregnant mice, even when abortion rates are increased following infusion of cytokines (where our data suggests OX2 continues to act to counter the effects of fgl2).

It is now known that OX2 functions following interaction with its receptor (OX2<sup>r</sup>) on target cells. At least 2 groups, ourselves (Gorczynski et al. 2000 and Wright et al. 2000) have documented the existence of OX2<sup>r</sup> on macrophages, and the inventors showed optimal inhibition of graft rejection in vivo occurred with infusion of both OX2:Fc and OX2r+ cells (Gorczynski et al. 2000). Unlike the Barclay group, the inventors have found that a large percentage (>80%) of ConA activated  $\gamma\delta$ T cells also express an OX2<sup>r</sup>, as defined by FACS with FITC-OX2:Fc. The inventors have recently confirmed this independently using our mAbs to OX2r, and following cDNA sequencing of the OX2r expressed in  $\gamma\delta$ TCR+ hybridomas (Yu et al: manuscript in preparation). The inventors have not yet studied the functional activity of



OX2r cells (whether macrophages or  $\gamma\delta$ T cells) in allopregnant mice. However, extrapolating from the data shown above, it is suggested that increased expression of OX2<sup>r</sup> will correlate with successful allopregnancy, and that triggering intracellular signaling by cross-linking OX2r on cells by mAb (presumably in the same fashion as native cell-bound OX2 does when it interacts with OX2<sup>r</sup>) rescues putatively doomed embryos mice from cytokine-induced spontaneous abortion (mediated by elevated fgl2 expression).

#### **EXAMPLE 9**

##### **OX-2 Expression Rescues Putatively Doomed Embryos**

The importance of CD200 (OX-2) in rescuing potentially doomed mouse embryos is illustrated by the in situ hybridization result in Figure 21. The pattern of hybridization with anti-sense probe for fgl2 (top panels) is compared with the pattern of hybridization using anti-sense probe for OX-2 for CBAxDBA/2 mated mice on day 8.5 of gestation. (Sense controls show no staining). Details of methods are provided in Molecular Human Reproduction 2001 (Clark et al. 2001). In section 1, left, fgl2 is ++ and OX-2 essentially negative. In section 2, fgl2 is still ++ but in this implantation, OX-2 hybridization can be seen in the same areas as fgl2. In section 3, fgl2 is only weakly expressed, and OX-2 is readily demonstrable. It has been shown that the observed rate of abortions is less than expected from the % of implantations showing fgl2<sup>++</sup>. When anti-OX-2 neutralizing antibody is administered, all of the fgl2<sup>++</sup> implantation appear to be aborted. This is attributed to the demise of implantations showing pattern #2.

#### **EXAMPLE 10**

##### **OX-2 is Expressed in Successful Human Pregnancy**

The importance of OX-2 (CD200) is rescue of potentially doomed fgl2<sup>hi</sup> mouse implantations has been shown above. Increased fgl2 expression is also seen in abortion of chromosomally normal human embryos in the first trimester, but not in chromosomally abnormal embryos. To determine if OX-2 is expressed in successful pregnancies (for which first trimester tissue is not available), term placentae were examined. The placentae were obtained after delivery, the decidua was scraped off, and the trophoblast cells were isolated by sequential enzymatic digestion, as

described by Guilbert et al. To remove non-trophoblast cells, anti-CD9 antibody was added, and the suspension was passed over an anti-immunoglobulin column, according to the method of Guilbert et al. The adhered CD9<sup>+</sup> 'stroma' was obtained by physical agitation of the beads of the column. RNA was extracted in the usual manner from a similar number of CD9<sup>+</sup> and CD9<sup>-</sup> cells and RT-PCR was performed (30 cycles) using primers to the full-length (3 exon) OX-2 gene, using primers to the 3rd exon that detects the truncated (exon 2-missing) mRNA. The PAGE gel in Figure 22 shows the molecular size ladder (lane 1), trophoblast with full length primers (lane 2) and exon 3 primers (lane 3). The full length and shorter mRNA OX-2 transcripts are seen. Lane 4 (corresponding to lane 2) and lane 5 (corresponding to lane 3) represent negative result obtained with CD9<sup>+</sup> stromal cells.

To test for OX-2 protein expression on trophoblast, the flow cytometry protocol (summarized at the end) was used with the CD9<sup>-</sup> fraction. Surface expression of OX-2 was detected using PE-tagged monoclonal antibody to human OX-2 (Cedarlane labs). Intracellular cytokeratin was stained using FITC-mAB from Dako. Figure 23 shows result using antibody detecting CK5,6,8,17; similar results have been obtained using FITC-anti-CK18. The upper left panel shows the forward and side scatter pattern. FITC and PE isotype controls used to set gates is shown below on left. Upper right panel shows cytokeratin-positive cells, the majority of which were also OX-2<sup>+</sup>. A significant population of CK- OX-2<sup>++</sup> cells was also noted. The identity of this population has not been established. On the right, the middle panel shows staining profile for OX-2 for the CK- and CK<sup>+</sup> populations, and the lower panel shows the scatter. Clearly these populations have different properties. Velocity sedimentation separation indicates both populations may have OX-2-dependent immunoregulatory activity.

#### **EXAMPLE 11**

##### **OX-2 Mab in the Prevention of Postcoital Pregnancy**

The inventors have shown that infusion of an immunoadhesin, OX-2:Fc, reduces the rate of abortion in a mouse model system using animals with increased rates of spontaneous abortion. Antibodies to OX-2 on the

other hand were shown to increase the rates of spontaneous abortion, and decrease litter size in this model. This allows the manipulation of fertility using non-hormonal methods.

5 This example confirms the ability of OX-2 Mab to reduce fertility in rodents.

The objection of this experiment is to determine if OX-2 Mab is capable of preventing or reducing the fertility in rodents after coitus.

Reduction in fertility is defined as:

- 10 a) the percent of treated impregnated females delivering pups is less than that observed in the control group.
- b) the average litter size of treated impregnated females carrying a litter to term is less than that observed in the control group.

### **Materials and Methods**

15 Rodents (rats/mice) were housed in individual cages with one female and one male/cage. Animals were inspected daily for vaginal plugs. Beginning on the day of fertilization (defined as the presence of a vaginal plug) females were kept one to a cage, and received ip injections of mAbs to OX2 rat anti-mouse (clone 3B6) for mice and mouse anti-rat (clone 6C2) for rats. The control group of the impregnated females received injections with control rat or mouse Ig.

20

Records were kept of the number of impregnated females producing litters in each group, as well as the litter size for each mother.

### **Animals**

Mouse female, male - outbred Swiss-Webster.

25 Rat female, male – outbred Sprague-Dawley.

### **Statistical Considerations**

Comparison of the percent of impregnated females carrying litters to term, and the mean litter size for successful pregnancies, were made independently for rats/mice using non parametric Chi-square or Fisher's Exact test where appropriate for pregnancy rates, and the Rank Sum test for litter sizes.

30

## **Methodology**

### **Breeding**

One animal of each sex is placed in a cage. The females are checked daily for vaginal plugs. Once a vaginal plug was observed the females were placed one to a cage.

Beginning on the day of impregnation (defined as the day a vaginal plug is observed), each female was moved to a new cage (housed alone) and begins treatment with mouse or rat OX2 antibodies by intraperitoneal injection.

### **Experiment No.1**

Mice: dose 100 micrograms per mouse every 3 days for 7 doses.

Rats: dose 300 micrograms per rat every 3 days for 7 doses

### **Experiment No.2**

Mice: dose 250 micrograms per mouse every 36 hours for up to 18 days

Rats: dose 500 micrograms per rat every 3 days for 7 doses

### **Methods of assessment**

Assessment of coitus: vaginal plug.

Assessment of pregnancy – number of pups delivered

In the second experiment the sex was also determined in each pup delivered.

The number of animals giving birth is counted as are the number of pups delivered.

Each female animal is housed in an individual cage until day 25 after breeding. Animals not delivering by day 25 were considered not to be pregnant and destroyed.

### **Study end points**

Number of animals in each group who delivered any pups.

Number of pups delivered to each animal achieving delivery at term.

### **Duration of the study**

Length of time the animal were held post impregnation....25 days.

Methods of termination....pups were sacrificed after recording the total litter size per female. In the second study the sex of each pup was also recorded.

Tissue to be taken...none.

### **Schedule of Evaluations**

Daily inspection of animals for impregnation (vaginal plugging)

Injection of pregnant animals every 3 days in the rat experiments and  
experiment 1 for the mice. The mice in experiment two were observed every  
5 36 hours.

Daily inspection from day 19 for evidence of birthing.

### **Test Article and Concomitant Treatments**

#### Test Article

Anti-OX2 mAbs were obtained from BioSpark, Canada.

10 Mouse anti-rat OX2: clone 6C2. Cat # SP 300X. Lot # 02

Stored at 1mg/ml (no azide) in PBS.

Rat anti-mouse OX2: clone 3B6. Cat # SP 200X Lot # 03

Stored at 1mg/ml, no azide, in PBS.

15 Control Ig was also obtained from BioSpark, Canada. This represents a  
preparation from pooled normal ascites (from rats or mice).

Again storage is in PBS, with no azide, at 1mg/ml.

Normal rat Ig: Lot #01. Cat. # SP89

Normal mouse Ig: Lot # 01; Cat # SP99

20

Inventory and accountability records are kept by the investigator

a) The investigator kept study drugs in a locked freezer, a secure storage  
facility, in his laboratory at the TGH (MBRW-2-926).

#### Experiment 1:

25 **Mice:**

Control: 17/17 mice impregnated had successful pregnancies.

Mean litter size: 7

3B6 injected: 17/18 mice impregnated had successful pregnancies.

Mean litter size: 6

30 See Table 8

Statistics: Mice: 17 control and 18 treated

The proportion of mice pregnant was not significantly different by Fisher's Exact test; the litter size in each group, and the average number of pups per group were not significantly different by the Rank Sum test.

**Rats:**

5 Control: 11/12 rats impregnated had successful pregnancies.

Mean litter size: 9

6C2 injected: 5/8 rats impregnated had successful pregnancies.

Mean litter size for the group was 5 but for the 5 rats delivering pups the mean litter size was 8

10 See Table 10

$P < 0.05$  for successful pregnancies.

Statistics: Rats 12 control and 8 treated

The proportion of animals pregnant in the treated group was not significantly reduced ( $P = 0.153$ , Fisher's Exact test). Litter size in those delivering was

15 reduced ( $P < 0.018$ , Rank Sum test), and the total number of pups generated by the immune treated group was reduced ( $P < 0.006$ , Rank Sum test)

**Experiment 2:**

**Mice:**

Control: 8/8 mice impregnated had successful pregnancies.

20 Mean litter size: 6

3B6 injected: 3/8 mice impregnated had successful pregnancies.

Mean litter size for the group was 2 but for the 3 rats delivering pups the mean litter size was 5

See Table 9

25 Statistics: Mice 8 control and 8 treated

The proportion of rats pregnant was significantly reduced by treatment ( $P = 0.0128$ , Fisher's Exact test), and the number of pups overall was reduced ( $P < 0.00021$ , Rank Sum test). The average litter size for animals successfully

30 delivering was also reduced ( $P < 0.024$ , Rank Sum test), but the number of litters was small.

**Rats:**

Control: 6/6 rats impregnated had successful pregnancies.

Mean litter size: 10

6C2 injected: 2/6 rats impregnated had successful pregnancies.

Mean litter size for the group was 2.5 but for the two rats delivering pups the mean litter size was 7.5

See Table 11

5 Statistics: Rats 6 control 6 treated

The proportion of animals that delivered litters was significantly reduced by treatment ( $P=0.03$ , Fisher's Exact test), as was the total number of pups produced by the treated group ( $P<0.002$ , Rank Sum test). Litter size was also slightly reduced in those delivering ( $P<0.03$ , Rank Sum test), but the number  
10 of litters was small.

### Conclusions

The administration of OX-2 antibodies to rats and mice following impregnations has been shown to reduce the number of animals delivering pups. The average liter size of antibody treated animals that did deliver was  
15 not different from the control animals.

While the present invention has been described with reference to what are presently considered to be the preferred examples, it is to be understood that the invention is not limited to the disclosed examples. To the  
20 contrary, the invention is intended to cover various modifications and equivalent arrangements included within the spirit and scope of the appended claims.

All publications, patents and patent applications are herein incorporated by reference in their entirety to the same extent as if each  
25 individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety.

**TABLE 1**

Summary of sequences and clones detected in cDNA library from pv immunized mice

5	<b>Match category</b>	<b>Number of clones represented (%)</b>
	Known mouse genes	30 (45)
	Non-mouse genes (rat/human) 10	14 (21)
	No data base match	22 (34)

15 **Footnotes:**

Genes were considered a “match” with a BLAST score >250 with a minimum of 50bp alignment.



**TABLE 2**

Cytokine production from cells of mice receiving pv immunization and anti-rat OX-2

Mabs given to recipients<sup>a</sup>

Cytokine levels in culture supernatants<sup>b</sup>

		IL-2	IFN $\gamma$	IL-4	IL-10
5	<b>No pv immunization (CsA only)</b>				
	<b>None</b>	750 $\pm$ 125	85 $\pm$ 18	29 $\pm$ 8	130 $\pm$ 40
	+anti-rat OX-2	890 $\pm$ 160	93 $\pm$ 19	30 $\pm$ 10	120 $\pm$ 35
	+anti-mouse CD28	415 $\pm$ 88*	57 $\pm$ 9*	105 $\pm$ 22*	275 $\pm$ 55*
10	+anti-mouseCTLA4	505 $\pm$ 125*	65 $\pm$ 8	95 $\pm$ 20*	190 $\pm$ 45
	+anti-B7-1	340 $\pm$ 65*	35 $\pm$ 7*	120 $\pm$ 21*	285 $\pm$ 60*
	+anti-B7-2	495 $\pm$ 90*	64 $\pm$ 7	90 $\pm$ 20*	185 $\pm$ 45
	<b>PV Immunization + CsA</b>				
15	<b>None</b>	190 $\pm$ 55	25 $\pm$ 8	107 $\pm$ 21	780 $\pm$ 150
	+anti-rat OX-2	730 $\pm$ 140*	60 $\pm$ 16*	33 $\pm$ 10*	220 $\pm$ 40*
	+anti-mouse CD28	145 $\pm$ 38	20 $\pm$ 9	145 $\pm$ 34	1140 $\pm$ 245
	+anti-mouseCTLA4	85 $\pm$ 25	15 $\pm$ 6	125 $\pm$ 31	960 $\pm$ 220
	+anti-B7-1	110 $\pm$ 30	20 $\pm$ 6	144 $\pm$ 28	885 $\pm$ 180
20	+anti-B7-2	75 $\pm$ 20	14 $\pm$ 5	150 $\pm$ 30	1230 $\pm$ 245

**Footnotes:**

a. 3 C3H mice/group were used in each experiment. All animals received CsA and C57BL/6 renal transplants as described in the Materials and Methods. Mice in the lower half of the Table also received pv infusions of  $15 \times 10^6$  C57BL/6 bone marrow derived dendritic cells on the day of transplantation. Where monoclonal antibodies were given the dose used was 100mg/mouse, x4 doses at 2 day intervals. All mice were sacrificed 14 days post transplantation. Spleen cells were cultured in triplicate from individual animals for 40hrs in a 1:1 mixture with irradiated C57BL/6 spleen stimulator cells.

b. Arithmetic mean ( $\pm$ SD) for triplicate determinations from individual samples of the animals described in the first column. All cytokines were assayed by ELISA. IL-2, IL-4 and IL-10 are shown as pg/ml, IFN $\gamma$  as ng/ml. Data are pooled from 2 such studies (total of 6 individual mice tested/group).

\* represents significantly different from control group with no Mab ( $p < 0.02$ )

**TABLE 3**

FACS staining of PBL and spleen adherent cells in different species, using anti-OX-2 Mabs

5	SPECIES <sup>a</sup>	Donor <sup>b</sup> Treatment	Mab	Percent stained cells <sup>c</sup>	
				PBL	Spleen
	Human	NONE	H4B4	1.5±0.3	4.8±1.7
			H4A9A2	1.5±0.4	6.1±2.0
			H4A9C7	1.3±0.4	4.3±1.7
10	Mouse	NONE	M3B5	1.9±0.4	6.7±2.1
			M3B6	1.7±0.4	5.2±1.6
			M2C8	1.4±0.4	4.2±1.4
15	Mouse	PV immune	M3B5	5.9±1.5	20±4.1
			M3B6	5.2±1.4	17±3.6
			M2C8	4.7±1.4	15±3.3
20	Rat	NONE	RC6A3	1.3±0.3	5.3±1.6
			RC6C2	1.5±0.4	6.5±1.7
			RC6D1	1.9±0.6	6.8±1.5
25	Rat	PV immune	RC6A3	4.8±1.3	16±4.2
			RC6C2	4.9±1.6	18±3.9
			RC6D1	5.3±1.7	20±4.5

**Footnotes:**

<sup>a</sup> Fresh cells were obtained from normal human donors (PBL), cadaveric transplant donors (human spleen), or from adult (8-10 week) mouse or rat donors. The same 3 separate tissue donors were used for each Mab tested.

<sup>b</sup> Donor pretreatment refers to infusion of allogeneic bone marrow cells into the portal vein (C57BL/6 for C3H mouse donors; BN for LEW rat donors) 4 days before harvest of PBL or spleen (see text and (6)).

<sup>c</sup> Arithmetic mean (+SD) for percent cells stained in 3 independent assays. Control antibodies (FITC anti-mouse IgG (for anti-human or anti-rat Mabs, or FITC anti-rat IgG for anti-mouse Mabs) gave no significant staining above background (<0.2%).

**TABLE 4**

Type-1 cytokine production in MLR cultures is increased by anti-OX-2 Mabs

Mabs in culture<sup>a</sup>

Cytokine levels in culture supernatants<sup>b</sup>

5		ELISA assays (murine only)				Bioassay (CTTL-2)	
		IL-2	IFN $\gamma$	IL-4	IL-10	IL-2	IL-6
	<b>MOUSE MLR</b>						
	None	350 $\pm$ 55	35 $\pm$ 18	345 $\pm$ 63	340 $\pm$ 50	480 $\pm$ 160	365 $\pm$ 74
10	M3B5	890 $\pm$ 160*	115 $\pm$ 29*	130 $\pm$ 10*	168 $\pm$ 42*	820 $\pm$ 200*	265 $\pm$ 46
	M3B6	915 $\pm$ 155*	117 $\pm$ 25*	135 $\pm$ 32*	135 $\pm$ 38*	850 $\pm$ 175*	303 $\pm$ 55
	M2C8	855 $\pm$ 155*	105 $\pm$ 28*	120 $\pm$ 32*	140 $\pm$ 37*	830 $\pm$ 165*	279 $\pm$ 61
	control Ig	370 $\pm$ 75	36 $\pm$ 11	330 $\pm$ 55	310 $\pm$ 45	335 $\pm$ 60	349 $\pm$ 59
	None**	710 $\pm$ 145	108 $\pm$ 23	110 $\pm$ 21	105 $\pm$ 23	690 $\pm$ 155	285 $\pm$ 54
15	<b>RAT MLR</b>						
	None					490 $\pm$ 145	360 $\pm$ 57
	RC6A3					690 $\pm$ 155*	295 $\pm$ 55
	RC6C2					845 $\pm$ 180*	345 $\pm$ 68
	RC6D1					830 $\pm$ 160*	370 $\pm$ 57
20	Control Ig					475 $\pm$ 160	356 $\pm$ 58
	<b>HUMAN MLR</b>						
	None					395 $\pm$ 85	295 $\pm$ 45
	H4B4					570 $\pm$ 125*	315 $\pm$ 50
	H4A9A2					630 $\pm$ 145*	320 $\pm$ 48
25	H4A9C7					625 $\pm$ 140*	345 $\pm$ 56
	Control Ig					360 $\pm$ 120	320 $\pm$ 50

Footnotes:

<sup>a</sup>. MLR cultures were set up as described in the Materials and Methods. For human MLR cultures the same 3 different responder preparations were used for each Mab, and stimulated with a pool of mitomycin C treated spleen stimulator cells (from a random mixture of 6 spleen donors). For mouse (C3H anti-C57BL/6) and rat (LEW anti-BN) MLR cultures all assays were set up in triplicate for each Mab. Mouse responder spleen cells were from mice treated 4 days earlier by portal vein infusion of C57BL/6 bone marrow cells, except for data shown as (None\*\*) where responder cells were from non-injected C3H mice. Mab was added as a 30% supernatant concentration. Supernatants were harvested for cytokine assays at 60 hrs.

<sup>b</sup>. Data show arithmetic means (+SD) for each Mab. For mouse assays all supernatants were assayed for a number of cytokines (ELISA), and for IL-2/IL-6 using bioassays (proliferation of CTLL-2, B9 respectively). Supernatants from rat/human cultures were assayed in bioassays

only. Note that cells incubated with isotype control Igs (non-reactive by ELISA or FACS) gave cytokine data indistinguishable from cultures incubated in the absence of Mab.  
 $p < 0.05$ , compared with cultures without Mabs.

Figure 1. Cytokine production by T cells in the presence of anti-CD3 and anti-CD28 antibodies. T cells were incubated with anti-CD3 (1 µg/ml) and anti-CD28 (1 µg/ml) for 48 hours. The supernatant was collected and analyzed for cytokine production by ELISA. The results are shown as mean ± SD of three independent experiments. \* indicates p < 0.05 compared to control.

**TABLE 5**

OX-2:Fc Immunoadhesin Inhibits Mixed Leukocyte Reaction *in vitro*

5	Added supernatant <sup>a</sup> (pg/ml) <sup>c</sup>	Percent lysis <sup>51</sup> Cr targets <sup>b</sup> (50:1, effector:target)	Cytokines in culture	
			IL-2	IL-4
	NONE (control)	31±4.0	1005±185	60±20
	Control CHO (vector transduced)	33±4.3	810±190	45±20
10	CHO transduced with OX-2:Fc	4.2±2.1	175±45	245±55

**Footnotes:**

15 <sup>a</sup> Supernatant was harvested at 7 days from CHO cells transduced with control pbK vector, or vector carrying a cDNA insert encoding OX-2 linked to murine Fc. A 1:1 mixture of supernatant was used in cultures containing 5x10<sup>6</sup> LEW spleen and 2.5x10<sup>6</sup> irradiated LBNF1 spleen cells; this corresponded to 50ng/ml OX-2:Fc

20 <sup>b</sup> and <sup>c</sup> Percent lysis with cells at 5 days, using 1x10<sup>4</sup> <sup>51</sup>Cr BN spleen ConA targets; cytokines in culture supernatants at 60hrs.

**TABLE 6**

Inhibition of skin graft rejection by OX-2:Fc

5	<b><u>Treatment of mice</u></b>	<b><u>Rejection of skin grafts (mean+SD) in days</u></b>
	NIL	12+3.8
	OX-2:Fc	19+4.2

10

Footnotes:

6 mice/group were treated as shown.

NIL indicates infusion of normal mouse IgG only.

Arithmetic mean (+SD) graft survival for group.

**TABLE 7**

OX-2:Fc infused into mice receiving skin allografts reverses polarization in cytokine production

	<b><u>Treatment of mice</u></b> <b><u>(pg/ml)</u></b>	<b><u>Cytokines in culture supernatant at 48hrs</u></b>	
		IL-2	IL-4
10	NIL	1250+160	80+20
	OX-2:Fc	350+85	245+50

**Footnotes:**

3 mice/group received iv infusions of saline or OX-2:Fc (50mg/mouse) every 2 days x4 from the time of grafting with C57BL/6 skin. Mice were sacrificed at 10 days and spleen cells stimulated *in vitro* with irradiated C57BL/6 spleen stimulator cells.

Arithmetic mean (+SD) for IL-2/IL-4 in supernatant at 48 hrs. Data are pooled from triplicate cultures for each mouse spleen.

**TABLE 8**

**Mice Experiment**

Mouse type: Swiss-Webster-Outbred

**Test Article:** Clone 3B6rat – anti-mouse OX-2

Dose 100 pg IP Q 3 days

**Placebo:** Mouse Igg

**Controls**

Mouse Number	Date Plugged	Delivery Date	Litter size		
1	28-Feb-01	Mar-21	8	Controls % Pregnant	100
2	Feb-29	Mar-21	6		
3	Feb-29	Mar-21	8		
4	Mar-01	Mar-22	6		
5	Mar-01	Mar-22	6		
6	Mar-03	Mar-24	8		
7	Mar-03	Mar-24	4		
8	Mar-05	Mar-26	5		
9	Mar-05	Mar-26	8		
10	Mar-06	Mar-27	8		
11	Mar-06	Mar-27	6		
12	Mar-06	Mar-28	8		
13	Mar-08	Mar-29	8		
14	Mar-08	Mar-29	6		
15	Mar-08	Mar-30	4		
16	Mar-08	Mar-30	8		
17	Mar-11	Mar-31	6		

**Treated**

1	Feb-28	Mar-20	7	Treated: % Pregnant	94
2	Feb-29	Mar-21	8		
3	Feb-29	Mar-21	8		
4	Feb-29	Mar-21	4		
5	Mar-01	Mar-22	5		
6	Mar-03	Mar-22	8		
7	Mar-03	Mar-23	8		
8	Mar-05	Mar-25	6		
9	Mar-05	Mar-25	6		
10	Mar-06	Mar-25	5		
11	Mar-06	Mar-26	4		
12	Mar-06	Mar-27	4		
13	Mar-06	Mar-27	8		
14	Mar-08	Nil	0		
15	Mar-08	Mar-29	6		
16	Mar-08	Mar-29	7		
17	Mar-08	Mar-29	8		
18	Mar-11	Mar-31	7		



**TABLE 9**

Mice Experiment

Mouse type: Swiss-Webster-Outbred

**Test Article:** Clone 3B6rat – anti-mouse OX-2

Dose 250 micrograms IP every 36 hours

**Placebo:** Mouse IgG

Mouse Number	Date Plugged	Delivery Date	Controls			
			Litter Size Males	Litter Size Females	Litter Size Total	
1	Apr-01	Apr-21	4	2	6	
2	Apr-01	Apr-22	4	3	7	
3	Apr-03	Apr-25	3	3	6	
4	Apr-05	Apr-26	2	4	6	
5	Apr-05	Apr-27	5	3	8	
6	Apr-08	Apr-29	4	4	8	
7	Apr-09	Apr-30	3	3	6	
8	Apr-10	Apr-30	2	4	6	

Mouse Number	Date Plugged	Delivery Date	Treated			
			Litter Size Males	Litter Size Females	Litter Size Total	
1	Apr-01	Nil	0	0	0	
2	Apr-02	Nil	0	0	0	
3	Apr-04	Apr-25	4	2	6	
4	Apr-05	Nil	0	0	0	
5	Apr-07	Apr-29	1	3	4	
6	Apr-09	Apr-30	4	2	6	
7	Apr-10	Nil	0	0	0	
8	Apr-13	Nil	0	0	0	

Controls    % Pregnant            100

Treated     % Pregnant            38

**TABLE 10**

**Rat Experiment**

**Rat type:** Sprague-Dawley-Outbred

**Test Article:** Clone 6C2 Mouse – anti-rat OX-2

**Dose** 300 micrograms IP Q 3 days

**Placebo:** Rat Igg

**Controls**

RAt Number	Date Plugged	Delivery Date	Litter size
1	Feb-29	Mar-21	11
2	Feb-29	Mar-22	13
3	Mar-01	Mar-24	10
4	Mar-01	Mar-24	11
5	Mar-03	Mar-25	12
6	Mar-03	Mar-25	9
7	Mar-05	Nil	0
8	Mar-05	Mar-27	9
9	Mar-06	Mar-28	8
10	Mar-07	Mar-28	9
11	Mar-07	Mar-29	10
12	Mar-09	Mar-31	8

Controls % Pregnant 92

**Treated**

Rat Number	Date Plugged	Delivery Date	Litter size
1	Feb-29	Mar-21	8
2	Mar-01	Mar-23	9
3	Mar-01	Nil	0
4	Mar-01	Mar-24	9
5	Mar-03	Nil	0
6	Mar-05	Nil	0
7	Mar-06	Mar-28	8
8	Mar-07	Mar-29	6

Treated: % Pregnant 63

**TABLE 11**

**Rat Experiment**

**Rat type:** Sprague-Dawley-Outbred

**Test Article:** Clone 6C2 Mouse – anti-rat OX-2

**Dose** 500 micrograms IP Q 3 days

**Placebo:** Rat Igg

**Controls**

Rat Number	Date Plugged	Delivery Date	Litter Size Males	Litter Size	Litter Size	Litter Size Total
				Females		
1	Apr-04	Apr-25	6	4		10
2	Apr-04	Apr-27	6	5		11
3	Apr-08	Apr-29	4	6		10
4	Apr-09	Apr-30	3	7		10
5	Apr-13	May-02	4	5		9
6	Apr-14	May-03	6	5		11

Controls % Pregnant 100

**Treated**

Rat Number	Date Plugged	Delivery Date	Litter Size Males	Litter Size	Litter Size	Litter Size Total
				Females		
1	Apr-05	Nil	0	0		0
2	Apr-06	Nil	0	0		0
3	Apr-09	Apr-30	3	4		7
4	Apr-12	Nil	0	0		0
5	Apr-14	Nil	0	0		0
6	Apr-16	May-06	4	4		8

Treated: % Pregnant 33

## FULL CITATIONS FOR REFERENCES REFERRED TO IN THE SPECIFICATION

- 5      **Ahvazi, B. C., P. Jacobs, and M. M. Stevenson.** 1995. Role of macrophage-derived nitric oxide in suppression of lymphocyte proliferation during blood-stage malaria. *J. Leu. Biol.* 58 (1):23-31.
- 10      **Akatsuka, Y., C. Cervený, and J. A. Hansen.** 1996. T cell receptor clonal diversity following allogeneic marrow grafting. *Hum. Immunol.* 48:125-134.
- 15      **Akolkar, P. N., B. Gulwani-Akolkar, R. Pergolizzi, R. D. Bigler, and J. Silver.** 1993. Influence of HLA genes on T cell receptor V segment frequencies and expression levels in peripheral blood lymphocytes. *J. Immunol.* 150 (April 1):2761-2773.
- 20      **Albina, J. E., J. A. Abate, and W. L. Henry.** 1991. Nitric oxide production is required for murine resident peritoneal macrophages to suppress mitogen-stimulated T cell proliferation. *J. Immunol.* 147:144-152.
- 25      **Banchereau, J., and R. M. Steinman.** 1998. Dendritic cells and the control of immunity. *Nature.* 392:245-252.
- 30      **Barclay, A.N.** 1981. Different reticular elements in rat lymphoid tissue identified by localization of Ia, Thy-1 and MRC OX-2 antigens. *Immunology* 44:727
- 35      **Barclay, A.N., and H.A. Ward.** 1982. Purification and chemical characterization of membrane glycoproteins from rat thymocytes and brain, recognized by monoclonal antibody MRC OX-2. *Eur. J. Biochem.* 129:447.
- Borriello, F., J. Lederer, S. Scott, and A.H. Sharpe.** 1997. MRC OX-2 defines a novel T cell costimulatory pathway. *J. Immunol.* 158:4548.
- Brasel, K., H. J. McKenna, K. Charrier, P. J. Morrissey, D. E. Williams, and S. D. Lyman.** 1997. Flt3 ligand synergizes with granulocyte-macrophage colony-stimulating factor or granulocyte colony-stimulating factor to mobilize hematopoietic progenitor cells into the peripheral blood of mice. *Blood.* 90:3781-3788.

- 5 **Bronstein, I., J.C. Voyta, O.J. Murphy, L. Bresnick, and L.J. Kricka.** 1992. Improved chemiluminescence Western blotting procedure. *Biotechniques* 12:748.
- Castle, B.E., K. Kishimoto, C. Stearns, M.L. Brown, and M.R. Kehry.** 1993. Regulation of the expression of the ligand for CD40 on T helper lymphocytes. *J. Immunol.* 151:1777.
- 10 **Chen, Z., H. Zeng, and R. M. Gorczynski.** 1997. Cloning and characterization of the murine homologue of the rat/human MRC OX-2 gene. *BBA. Mol. Basis Dis.* 1362:6-10.
- 15 **Chen, Z. and Gorczynski, R.M.** 1997. *Biochem. Biophys. Acta.* 100, in press.
- 20 **Clark, D.A., J.-W. Ding, G. Yu, G.A. Levy and R. Gorczynski.** 2001. Fgl2 prothrombinase expression in mouse trophoblast and decidua triggers abortion but may be counted by OX-2. *Mol. Human Reproduction*, Vol. 7, No. 2, pp. 185-194.
- 25 **Freeman, G. J., V. A. Boussiotis, A. Anumanthan, G. M. Bernstein, X. Y. Ke, P. D. Rennert, G. S. Gray, J. G. Gribben, and L. M. Nadler.** 1995. B7-1 and B7-2 do not deliver identical costimulatory signals, since B7-2 but not B7-1 preferentially costimulates the initial production of IL-4. *Immunity.* 2:523-532.
- 30 **Garside, P., and A. M. Mowat.** 1997. Mechanisms of oral tolerance. *Crit. Rev. Immunol.* 17:119-137.
- Goodwin, R.G., Din, W.S., Davis-Smith, T. et al.** 1993. *Eur. J. Immunol.* 23, 2631-2641
- 35 **Gorczynski, R. M.** 1992. Immunosuppression induced by hepatic portal venous immunization spares reactivity in IL-4 producing T lymphocytes. *Immunol. Lett.* 33:67-77.

**Gorczynski, R. M., and D. Wojcik.** 1992. Antigen presentation by murine splenic, but not hepatic, antigen-presenting cells to induce IL-2/IL-4 production from immune T cells is regulated by interactions between LFA-1/ICAM-1. *Immunol. Lett.* 34:177-182.

5

**Gorczynski, R. M., and D. Wojcik.** 1994. A role for non-specific (cyclosporin A) or specific (monoclonal antibodies to ICAM-1, LFA-1 and interleukin-10) immunomodulation in the prolongation of skin allografts after antigen-specific pre-transplant immunization or transfusion. *J. Immunol.* 152:2011-2019.

10

**Gorczynski, R.M., Z. Chen, S. Chung, Z. Cohen, G. Levy, B. Sullivan, and X.-M. Fu.** 1994a. Prolongation of rat small bowel or renal allograft survival by pretransplant transfusion and/or by varying the route of allograft venous drainage. *Transplantation* 58:816-820.

15

**Gorczynski, R.M.** 1994b. Adoptive transfer of unresponsiveness to allogenic skin grafts with hepatic  $\gamma\delta$ + T cells. *Immunology* 81:27-35.

20

**Gorczynski, R. M.** 1995a. Regulation of IFN $\gamma$  and IL-10 synthesis *in vivo*, as well as continuous antigen exposure, is associated with tolerance to murine skin allografts. *Cell. Immunol.* 160:224-231.

25

**Gorczynski, R. M., N. Hozumi, S. W. Wolfe, and Z. Chen.** 1995b. Interleukin-12, in combination with anti-interleukin-10, reverses graft prolongation after portal venous immunization. *Transplantation.* 60:1337-1341.

30

**Gorczynski, R.M., Z. Cohen, X.M. Fu, Z. Hua, Y.L. Sun, and Z.Q. Chen.** 1996a. Interleukin-13, in combination with anti interleukin-12, increases graft prolongation after portal venous immunization with cultured allogenic bone marrow-derived dendritic cells. *Transplantation* 62:1592-1600.

35

**Gorczynski, R. M., Z. Cohen, G. Levy, and X. M. Fu.** 1996b. A role for gamma delta TCR(+) cells in regulation of rejection of small intestinal allografts in rats. *Transplantation.* 62:844-851.

**Gorczynski, R. M., Z. Chen, Y. Hoang, and B. RossiBergman.** 1996c. A subset of gamma delta T-cell receptor-positive cells produce T-helper type-2

cytokines and regulate mouse skin graft rejection following portal venous pretransplant preimmunization. Immunology. 87 (3):381-389.

5 **Gorczynski, R. M., Z. Chen, H. Zeng, and X. M. Fu.** 1998a. A role for persisting antigen, antigen presentation and ICAM-1 in the increased renal graft survival following oral or portal vein donor-specific immunization. Transplantation. 66: 000-008.

10 **Gorczynski, R. M., Z. Chen, X. M. Fu, and H. Zeng.** 1998b. Increased expression of the novel molecule Ox-2 is involved in prolongation of murine renal allograft survival. Transplantation. 65:1106-1114.

15 **Gorczynski, R.M., et al.** 1998c. Analysis of cytokine production and V beta T-cell receptor subsets in irradiated recipients receiving portal or peripheral venous reconstitution with allogeneic bone marrow cells, with or without additional anti-cytokine monoclonal antibodies. Immunology. 93: p. 221-229.

20 **Gorczynski, R.M., Chen, Z., Fu, X.M. and Zeng, H.** 1998. J. Immunol. 160, in press

**Gorczynski, R.M., Chen, Z., Zeng, H. and Fu, X.M.** 1998. Transplantation submitted

25 **Gruss, H. and Dower, S.** 1995. Blood 85, 3378-3404

**Hancock, W. W., M. H. Sayegh, X. G. Zheng, R. Peach, P. S. Linsley, and L. A. Turka.** 1996. Costimulatory function and expression of CD40 ligand, CD80, and CD86 in vascularized murine cardiac allograft rejection. Proc. Natl. Acad. Sci. USA. 93:13967-13972.

30 **Jenkins, M. K., J. D. Ashwell, and R. H. Schwartz.** 1988. Allogeneic non-T spleen cells restore the responsiveness of normal T cell clones stimulated with antigen and chemically modified antigen-presenting cells. J. Immunol. 140:3324-3329.

35 **Kenick, S., R. P. Lowry, R. D. S. Forbes, and R. Lisbona.** 1987. Prolonged cardiac allograft survival following portal venous inoculation of allogeneic cells: What is "hepatic tolerance?". Transpl. Proc. 19:478-480.

- Kohler, G. and C. Milstein.** 1975. Preparation of monoclonal antibodies. *Nature*. 25: p. 256-259.
- 5     **Kronin, V., K. Winkel, G. Suss, A. Kelso, W. Heath, J. Kirberg, H. vonBoehmer, and K. Shortman.** 1996. Subclass of dendritic cells regulates the response of naive CD8 T cells by limiting their IL-2 production. *J. Immunol.* 157:3819-3827.
- 10    **Kuchroo, V. K., M. P. Das, J. A. Brown, A. M. Ranger, S. S. Zamvil, A. Sobel, H. L. Weiner, N. Nabavi, and L. H. Glimcher.** 1995. B7-1 and B7-2 costimulatory molecules activate differentially the Th1/Th2 developmental pathways: application to autoimmune disease therapy. *Cell*. 80:707-718.
- 15    **Larsen, C. P., S. C. Ritchie, R. Hendrix, P. S. Linsley, K. S. Hathcock, R. J. Hodes, R. P. Lowry, and T. C. Pearson.** 1994. Regulation of immunostimulatory function and costimulatory molecule (B7-1 and B7-2) expression on murine dendritic cells. *J. Immunol.* 152:5208-5219.
- 20    **Larsen, C.P., Elwood, E.T., Alexander, D.Z. et al.** 1996. *Nature* 381, 434-438
- 25    **Larsen, C. P., and T. C. Pearson.** 1997. The CD40 pathway in allograft rejection, acceptance, and tolerance. *Curr. Opin. Immunol.* 9:641-647.
- 30    **Leenen, P. J. M., K. Radosevic, J. S. A. Voerman, B. Salomon, N. vanRooijen, D. Klatzmann, and W. vanEwijk.** 1998. Heterogeneity of mouse spleen dendritic cells: *In vivo* phagocytic activity, expression of macrophage markers, and subpopulation turnover. *J. Immunol.* 160:2166-2173.
- 35    **Lenschow, D. J., T. L. Walunas, and J. A. Bluestone.** 1996. CD28/B7 system of T cell costimulation. *Annu. Rev. Immunol.* 14:233-258.
- Maraskovsky, E., K. Brasel, M. Teepe, E. R. Roux, S. D. Lyman, K. Shortman, and H. J. McKenna.** 1996. Dramatic increase in the numbers of functionally mature dendritic cells in Flt 3 Ligand-treated mice: multiple dendritic cell subpopulations identified. *J. Exptl. Med.* 184:1953-1962.



**Mayer, L.** 1996. Yin and Yang of mucosal immunology. *Transpl. Proc.* 28:2435-2437.

- 5     **McCaughan, G.W., et al.** 1987. The gene for MRC OX-2 membrane glycoprotein is localized on human chromosome 3. *Immunogenetics.* 25: p. 133-135.

- 10    **Miller, R. G., and R. A. Phillips.** 1969. The separation of cells by velocity sedimentation. *J. Cell. Comp. Physiol.* 73:191-198.

**Preston, S., et al.** 1997. The leukocyte/neuron cell surface antigen OX2 binds to a ligand on macrophages. *Eur J Immunol.* 27(8): p. 1911-8.

- 15    **Qian, J. H., T. Hashimoto, H. Fujiwara, and T. Hamaoka.** 1985. Studies on the induction of tolerance to alloantigens. I. The abrogation of potentials for delayed-type hypersensitivity responses to alloantigens by portal venous inoculation with allogeneic cells. *J. Immunol.* 134:3656-3663.

- 20    **Rosenzwajg, M., S. Camus, M. Guigon, and J. C. Gluckman.** 1998. The influence of interleukin (IL)-4, IL-13, and Flt3 ligand on human dendritic cell differentiation from cord blood CD34(+) progenitor cells. *Exp. Hematol.* 26:63-72.

- 25    **Salomon, B., J. L. Cohen, C. Masurier, and D. Klatzmann.** 1998. Three populations of mouse lymph node dendritic cells with different origins and dynamics. *J. Immunol.* 160:708-717.

- 30    **Sandhu, G.S., B.W. Eckloff, and B.C. Kline.** 1991. Chemiluminescent substrates increase sensitivity of antigen detection in Western blots. *Biotechniques* 11:14.

**Schwartz, R.H.** 1996. Models of T cell anergy: Is there a common molecular mechanism? *J Exp Med.* 184: p. 1-8.

- 35    **Steinbrink, K., M. Wolfl, H. Jonuleit, J. Knop, and A. H. Enk.** 1997. Induction of tolerance by IL-10-treated dendritic cells. *J Immunol.* 159:4772-4780.

- 5     **Steptoe, R. J., F. M. Fu, W. Li, M. L. Drakes, L. A. Lu, A. J. Demetris, S. G. Qian, H. J. McKenna, and A. W. Thomson.** 1997. Augmentation of dendritic cells in murine organ donors by Flt3 ligand alters the balance between transplant tolerance and immunity. *J Immunol.* 159:5483-5491.
- 10    **Suss, G., and K. Shortman.** 1996. A subclass of dendritic cells kills CD4+ T cells via Fas/Fas-ligand-induced apoptosis. *J. Exptl. Med.* 183:1789-1796.
- 15    **Swain, S. L.** 1995. Who does the polarizing? *Curr. Biol.* 5:849-851.
- Thelen, M., and U. Wirthmueller.** 1994. Phospholipases and protein kinases during phagocytic activation. *Curr. Opin. Immunol.* 6:106-112.
- Zheng, X.X., Steele, A.W., Nickerson, P.W. et al.** 1995. *Journal of Immunology* 154, 5590-5600.